



HAL
open science

Microplastic inputs to the Mediterranean Sea during wet and dry seasons: The case of two Lebanese coastal outlets

Rosa Sawan, Périne Doyen, Florence Viudes, Rachid Amara, Céline Mahfouz

► **To cite this version:**

Rosa Sawan, Périne Doyen, Florence Viudes, Rachid Amara, Céline Mahfouz. Microplastic inputs to the Mediterranean Sea during wet and dry seasons: The case of two Lebanese coastal outlets. *Marine Pollution Bulletin*, 2024, 198, pp.115781. 10.1016/j.marpolbul.2023.115781 . hal-04394933

HAL Id: hal-04394933

<https://ulco.hal.science/hal-04394933v1>

Submitted on 9 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

26 I. Introduction

27 Increased plastic production and consumption, illegal waste disposal and lack of waste management are
28 considered the main causes of plastics in the environment worldwide (Li et al., 2018). Microplastics (MPs)
29 ranging in dimension from 1 μm to 5 mm are considered a widespread and omnipresent contaminant
30 (Zhang et al., 2019b). They have been detected in various environmental compartments, namely marine
31 and inland water such as lakes and rivers (Auta et al., 2017; Dris et al., 2015a; van Emmerik et al., 2018;
32 Rios et al., 2019; Sarijan et al., 2021; Talbot and Chang 2022), sediments (Adomat and Grischek 2021;
33 Kundu et al., 2022) and atmospheric compartments (Zhang et al., 2020). Their accumulation in ecosystems
34 presents a relevant toxicological risk, as they can adsorb pollutants on their surfaces, and expose biota
35 through direct and indirect consumption (Constant et al., 2020; de Carvalho et al., 2021; Park, 2021).
36 Besides, MPs have already been found in many aquatic organisms such as bivalves, fish, and crustaceans
37 (Sarijan et al., 2021).

38 Among the world's basins, the semi-enclosed Mediterranean Sea Basin, shared by 21 countries, is one of
39 the most studied maritime areas in terms of MP pollution, with, for example, studies on sea surface water
40 (Panti et al., 2015; Ruiz-Orejón et al., 2016; Cincinelli et al., 2019), seafloor and coastal sediments (Alomar
41 et al., 2016; Abidli et al., 2018; Kazour et al., 2019; Fagiano et al., 2023;), as well as freshwater discharged
42 into the basin (Rayadin et al., 2016; Campanale, et al., 2020; Constant et al., 2020). However, the
43 occurrence of MPs in these freshwater systems has generally been less well studied and understood than
44 in marine systems (Rodrigues et al., 2018; Wong et al., 2020b; Treilles et al., 2021b). The first conclusions
45 drawn by studies in freshwater are that MP contamination depends on different variables, such as land
46 use, urban growth, hydrological conditions, physical characteristics of rivers, and agricultural activities (de
47 Carvalho et al., 2021; Wang et al., 2021b). Stormwater runoff has been identified as a major source of
48 plastic pollution in rivers, as it is the main mobilizer and a vector of terrestrial microplastics (Piñon-Colin
49 et al., 2020; Shruti et al., 2021). Moreover, freshwater discharged from rivers and sub-basins plays a crucial
50 role in the dynamic transportation of MPs from terrestrial to aquatic environments, with, for example,
51 different sources such as sewage discharge, recreational activities and wind strength (Piñon-Colin et al.,
52 2020). Recently, the impact of meteorological conditions on the fate and transportation of MPs into rivers
53 has also been investigated, with a certain positive correlation reported between precipitation and the
54 abundance of MPs (Constant et al., 2020; Xia et al., 2020), while others have reported a negative
55 correlation (Balthazar-Silva et al., 2020; Wicaksono et al., 2021).

56 In Lebanon, located along the eastern shore of the Mediterranean Sea, no study has assessed freshwater
57 contamination by MPs. Most studies on freshwater quality have, in fact, focused on physico-chemical or
58 geochemical composition to address the impact of poor wastewater management, landfill sites and
59 population growth (Falah et al., 2021; Maatouk et al., 2022). Only a few studies have been conducted
60 along the Lebanese coast to investigate the MP contamination of coastal surface sea water, biota and
61 marine sediments (Kazour et al., 2019; Jemaa et al., 2021; Mahfouz et al., 2023). This study aims to provide
62 an overview of the presence, contamination and variability of microplastics in two coastal Lebanese rivers.
63 The Ibrahim River and the Antelias River were chosen with a view to defining their respective inputs of
64 MPs to the Mediterranean Sea, focusing on (1) the impact of land use and urbanization around these
65 rivers on the abundance and type of MPs, and (2) the impact of two different meteorological conditions:
66 dry and wet seasons, on this plastic pollution. This will provide a better understanding of the sources of
67 pollution and will make it possible to quantify and qualify the flow of microplastic pollution into the
68 Mediterranean Sea. The data collected will form a crucial baseline source of information to assist with the
69 development of a strategic approach to address plastic pollution in aquatic ecosystems.

70

71 II. Material and methods

72 1. Study sites

73 Two river outlets were chosen prior to the mixing zone between seawater and freshwater from two less
74 or more urbanized watersheds: the Ibrahim River outlet (Site 1, S1) and the Antelias River outlet (Site 2,
75 S2) (**Figure 1**).

76 The Ibrahim Basin (**Figure 1**; S1) is located in the west-central region of Lebanon, 20 km from Beirut and
77 it drains directly into the Mediterranean Sea (El Najjar et al., 2019). It is a karstic watershed with distinct
78 hydrological systems and flows, making it a complex and challenging area of study. It takes its source from
79 two main springs, Afqa and Roueiss, at altitudes of 1,200 m and 1,300 m respectively (Assaker 2016). The
80 mean annual flow of the Ibrahim River ranges between 10.70 and 12.77 m³/s (Hanna et al., 2018). It is
81 one of the largest in Lebanon, with a catchment area of 330 km², a river length of 28 kms and a population
82 density of 1,700 inhabitants/km² (Daou et al., 2013; El Najjar et al., 2020). This watershed is occupied at
83 95% by natural and semi-natural zones and at 5% by urban and industrial areas, home to small businesses
84 such as food establishments and other plastic and paint manufacturing plants (Daou et al., 2013; Hanna
85 et al., 2018). Untreated domestic sewage and industrial effluents are discharged directly into the river due
86 to the lack of sewerage systems and wastewater treatment facilities (Slim et al., 2005; Hanna et al., 2018).

87 The Antelias Basin (**Figure 1**; S2) is located 11 km north of Beirut and represents an area of 24 km². It has
88 a population density of 19,486 inhabitants/km² and drains directly into the Mediterranean Sea (Nehme et
89 al. 2013). The Antelias River has an underground mountain river as a source (Hakim et al., 1988), while
90 82% of the population is connected to a combined sewerage system that discharges directly into the sea
91 through three outlets. However, the main river course receives raw sewage from some city buildings and
92 industries (Maatouk 2014). This river is heavily contaminated by untreated sewage, as shown by the high
93 biological oxygen demand (BOD), and the high concentrations of E.coli and coliform (Hourri et al, 2007).
94 38% of this watershed's land is occupied by urban and industrial areas, including for the production of
95 textiles, PVC and printing (Maatouk, 2014).

96 2. Sampling methodology and laboratory treatment

97 A series of seven sampling campaigns were conducted on the river outlets, the Ibrahim River (S1) and the
98 Antelias River (S2) between September 2021 and January 2022 (**Supplementary data, Table 1**). Three
99 sampling campaigns were performed during the dry season, with no rainfall over the last ten days,
100 whereas four sampling campaigns were conducted during the wet season, with cumulative precipitation
101 ranging from 16.8 mm to 88.8 mm (**Supplementary data, Table 1**). Lebanon has a Mediterranean climate,
102 with annual precipitation during the period from 1991 to 2021 averaging between 700 mm and 1,000 mm
103 along the coast. In January and December, annual precipitation averaged 157.34 mm and 135.99 mm
104 respectively. It is noteworthy that approximately 70% of this rainfall is typically concentrated within a few
105 days (the World Bank, 2021). During our study, an overall total of 307.2 mm was recorded during the rainy
106 season with 25 days of rain. A 100 L sample of surface water was collected at each outlet using a 10 L
107 metal bucket, thrown 10 times towards the center of the stream. The collected water was filtered through
108 a series of two stainless steel sieves with a mesh size of 500 µm and 150 µm, respectively. The sieves were
109 rinsed with filtered, distilled water into clean glass bottles to recover the particles. At the laboratory, the
110 recovered samples were placed at 40°C until the water had evaporated completely. 500 ml of 10% (w/v)
111 KOH solution (Chimieplus, Saint-Paul-de-Varax, France) was then added to each dried bottle to digest the
112 organic matter. Samples were stirred at 300 rpm and incubated at 40°C for 48 hrs on a magnet-heated
113 plate. The digested solution obtained was filtered on GF/A glass fiber filters with a porosity of 1.6 µm
114 (Cytiva, Freiburg im Breisgau, France). The filters were then carefully stored in glass Petri dishes at room
115 temperature until the observation and identification of particles.

116 3. Microparticle observation and identification

117 Filters were observed with a Leica M165 C stereoscope under two magnifications (x80 and x100). A zigzag
118 observation method was used, which involved observing the filter from top to bottom and then from

119 bottom to top, moving to the right. Pictures of suspected MPs were taken with a Leica M170 HD camera
120 and LAS (Leica Application Suite) software. All suspected MPs were counted, categorized by type (films,
121 fragments, microbeads or fibers) and color. The size of each suspected item was measured using
122 "Piximètre" software version 5.9 and four size categories were defined: [150 μm – 500 μm], [500 μm –
123 1,000 μm], [1,000 μm – 3,000 μm] and [3,000 μm – 5,000 μm].

124 After visual observation, the polymer nature of the suspect particles was identified using the Micro-Raman
125 Spectrometer XploRA™ Plus (HORIBA Scientific®, France), with two lasers (532 nm and 785 nm), a spatial
126 resolution ranging from 1 mm to 0.9 μm , and a spectrum ranging from 200 to 3,400 cm^{-1} . This device is
127 equipped with a self-contained box preventing airborne contamination. Each spectrum of particles was
128 compared to a polymer database identification (KnowItAll, WILEY®) and a personal library compiled using
129 standard polymers obtained from Goodfellow (France). Polymer identification via Raman microscopy
130 followed the procedure mentioned in Kazour et al. (2019). $\mu\text{FT-IR}$ was used when certain particles were
131 too dark to be identified by the Micro-Raman due to their color. A particle concentration, expressed as
132 particles/L, was calculated for each sample.

133 4. Quality Assurance and Control (QA/QC)

134 Numerous precautions to prevent plastic contamination were carried out throughout this study. Glass and
135 stainless steel materials only were used (Petri dishes, Erlenmeyer flasks, a filtration system, etc.). During
136 laboratory procedures, cotton lab coats and nitrile gloves were worn. Each solution used during analyses,
137 Milli-Q® water (Synergy UV), 70% (v/v) ethanol, 10% (w/v) KOH) was filtered on GF/A glass fiber filters (1.6
138 μm pore size). Milli-Q® water and filtered 70% ethanol were used to rinse all the equipment. All procedures
139 were carried out under a Thermo Scientific™ Herasafe™ 2030i laminar flow cabinet (Saint-Herblain,
140 France) in order to prevent atmospheric contamination of the samples. Before starting the analyses, the
141 bench of the laminar flow cabinet was cleaned with Milli-Q water, 70% (v/v) ethanol solution and Milli-Q®
142 water again.

143 Additionally, a set of different controls was used to control contamination. A field blank, consisting of 25
144 ml of filtered distilled water placed in an open container beside us on the banks of the river was performed
145 during each sampling campaign in order to consider potential atmospheric microplastic contamination.
146 These field blanks were filtered to count particles. Negative controls, i.e. an Erlenmeyer flask containing
147 500 mL of 10% KOH without a sample was processed and analyzed at the same time as the samples to
148 check that there was no contamination during handling. Additional atmospheric controls, i.e. GF/A glass

149 fiber filters placed into glass Petri dishes, were carried out during the digestion and filtration steps in order
150 to monitor the level of contamination inside the laminar flow cabinet.

151 The particles observed on all controls were counted following the size range in this study (between 150
152 μm and 5,000 μm) and deducted from the sample filters according to shape, size, and color.

153 5. Environmental data

154 The daily precipitation data were obtained from the official meteorological services "Meteo Liban". Land
155 cover-use shapefile data were provided by the National Council for Scientific Research in Lebanon (CNRS)
156 and the daily flow rate data were obtained from the official authority, Litani River Authority (LRA). These
157 data were expressed in two forms: D for daily precipitation of the sampling day and D-10, representing
158 cumulative precipitation during the 10 days prior to sampling plus rainfall on the day of sampling.

159 6. Data treatment

160 Statistical analyses were carried out using SPSS software (IBM SPSS STATISTICS 26) and R software. Since
161 the assumptions of normality and homogeneity of variance required for parametric statistics were not
162 met, the Kruskal-Wallis and Mann-Whitney tests were used to compare the variation of MPs in terms of
163 abundance, size and the nature of polymers, between Sites S1 and S2, as well as to compare the temporal
164 variation in MPs between the sites. Finally, correlations between MP abundance, types, and
165 environmental parameters such as daily precipitation, cumulative precipitation and river flow rate were
166 tested using the Spearman's rank correlation coefficient. The level of significance was set at $\alpha=0.05$.

167 III. Results and discussion

168 1. Impact of urbanization

169 a) The abundance of particles in the rivers studied

170 Microplastics were observed in all water samples with concentrations ranging between 0.58 and 4.59
171 particles/L at Site 1 (the Ibrahim River outlet), and between 2.2 and 28.91 particles/L at Site 2 (the Antelias
172 River outlet) (**Figure 2, a**). Site 2 exhibited significantly higher levels of particles (14.04 ± 9.8 particles/L),
173 compared to Site 1 (1.74 ± 1.38 particles/L) (Mann-Whitney Test, $p < 0.05$).

174 The levels of particles measured in this study are higher compared to those recorded in other
175 Mediterranean tributaries. The abundance of particles in the largest river of southeast Italy ranged from
176 0.9 ± 0.4 to 13 ± 5 particles/L and eight rivers sampled at Mersin Basin, Turkey (north-eastern part of the
177 Mediterranean Basin) revealed concentrations between 0.29 ± 0.06 particles/L (Campanale et al., 2020;
178 Demir, 2022). Two other Mediterranean rivers, the Rhône River and the Têt River, were also less
179 contaminated, with concentrations ranging between 0.0003 and 0.6185 items/L (Constant et al., 2020).

180 In addition, the reported MP concentrations were relatively higher than those reported at the Ebro River
181 in the north-western Mediterranean Basin, 0.0035 ± 0.0014 items/L (Simon-Sánchez et al. 2019). The
182 Yangtze River in China also exhibited lower concentrations (0 to 0.259 particles/L) compared to our values
183 (Li et al., 2020). However, concentrations reported in this study showed similar levels compared to a
184 sophisticated Shanghai river system with a concentration between 0.7 and 24.3 items/L (Wang et al.,
185 2021b) and the surface waters of urban rivers in Chengdu, the largest city in western China (5.00 – 10.5
186 items/L) (Chen et al., 2022).

187 However, the comparison with these reported MP concentrations is complicated due to the different
188 sampling techniques and sample processes used. These different studies cited the respective use of a
189 plankton net, a 2 L beaker, a manta trawl, a metal cylinder and a water sampler to collect different
190 volumes of water ranging from 10 L to 500 m³ (Campanale et al., 2020; Constant et al., 2020; Demir 2022;
191 Li et al., 2020; Wang et al., 2021b). These different sampling methods as well as the lower limit of the MP
192 analyzed, which depends on the mesh size of the analytical technique used, may result in varying degrees
193 of abundance and differing characterization of the microplastics. Many reviews have highlighted the
194 importance of protocol standardization to evaluate the presence of microplastic pollution in the
195 environment (Horton et al., 2017; Li et al., 2018; Pasquier et al., 2022).

196 Despite the impossibility of carrying out a direct comparison between studies, it is interesting to note that
197 the trend observed in this study was in line with other studies on the degree of urbanization. The highest
198 abundance of particles was recorded at Site 2, where 38% of its surface area is occupied by urban and
199 industrial zones, compared to Site 1, which covers only 5%. This correlation between abundance and
200 urbanization has already been observed in three estuaries on the east coast of Australia where the particle
201 concentration increased as the level of urbanization increased (Hitchcock et al., 2019). Moreover, Site 2
202 is defined as a very high traffic area due to its highly anthropogenic activities, leading to the dumping of
203 waste such as bottles, cigarette butts and other plastics directly onto the banks and consequently into the
204 river. The abundance measurements obtained in this study supported the link between MP pollution in
205 rivers and the degree of urbanization, which is coherent with previous studies highlighting the significant
206 contribution of the urbanization to MP pollution of river ecosystems (Wu et al., 2019; Constant et al.,
207 2020; Wong et al., 2020a; Piñon-Colin et al., 2020; Stovall & Bratton, 2022; Kunz et al., 2023).

208 The significant level of contamination reported in this study can be attributed to mismanagement of solid
209 and liquid waste near the rivers, where it is estimated that up to 90% of all wastewater (urban and
210 industrial) in Lebanon is discharged directly into its rivers, lakes or the sea without treatment, leading to

211 environmental and health degradation (Khatib et al., 2018). Moreover, open and closed landfills are
212 located near the aquatic environment (Kazour et al., 2019).

213 b) Morphological and polymeric variation of MPs across sites

214 All shapes of MPs (**Supplementary data, Figure 1**) were observed at both sites. There was a predominance
215 of fibers followed by fragments, films and finally microbeads (**Figure 2, b**). The more urbanized Site 2 had
216 a significantly higher percentage of fibers (88%) than Site 1 (72%) (Kruskal-Wallis, p-value < 0.05), which
217 could be related to a release due to the degradation of clothing, textile blankets and other products,
218 notably through washing machines used by residents (Napper et al., 2016; Hitchcock et al., 2019). These
219 results are consistent with other studies showing a predominance of fibers in urban rivers (Dris et al.,
220 2015b; Wang et al., 2017; Luo et al., 2019). Furthermore, S2 showed a significantly higher percentage of
221 films than S1, probably originating from the fragmentation of plastic packs, such as bottles, plastic wraps
222 and bags (Rowley et al. 2020), which would not be unexpected given the high level of human activity along
223 the Antelias River.

224 The particle size ranged from 150 μm to 4930 μm at Site S1 and 150 μm to 4604 μm at Site S2. Our findings
225 show significant differences in particle size between the two sites (Mann-Whitney, p-value < 0.05). The
226 size of plastic particles is also often linked to physical conditions, such as exposure time to the sun and UV
227 light, or the physical conditions at the different sites, including wind action, sediments. These conditions
228 can mechanically break down large plastic objects into smaller particles and, lastly, could be linked to
229 biological activities such as the formation of biofilms (Zhang et al. 2021). This difference could be linked
230 to the source of pollution, as S2 is under pressure from urbanization and anthropogenic activities. With
231 regard to S1, the river has an annual flow with a high mean annual discharge of $15 \pm 17 \text{ m}^3/\text{s}$ compared to
232 S2 ($0.7 \pm 0.73 \text{ m}^3/\text{s}$) which influence the mechanical breakdown and the formation of biofilms. Most of the
233 fibers were smaller than 1,000 μm throughout the study, (**Figure 2, c**). Our results are consistent with
234 several studies where microplastics between 100 μm and 1,000 μm were more frequently observed than
235 other size ranges (He et al., 2021; Luo et al., 2019; Su et al., 2020; Treilles et al., 2021b).

236 Regarding colors, blue and black particles were predominant across the samples, representing 40% and
237 66% of the particles at S1 and S2 respectively (**Figure 2, d**). These observations were in line with a recent
238 study carried out in coastal areas of Lebanon where blue-colored items had the highest concentrations,
239 followed by white and black items (Jemaa et al., 2021). The diversity of colors observed in this study
240 suggests a diversity of particle sources (Zhang et al., 2019a) and may represent a hazard, as colored
241 microplastics can sometimes be confused as food and ingested by aquatic fauna (Lopes et al., 2020).

242

243 A total of 551 particles corresponding to 5% of the total observed particles were analyzed using a Micro
244 Raman and a μ -FTIR (n=15) in order to identify their nature (**Figure 3, a; Supplementary data, Table 2**).

245 71.8% of these analyzed particles (n=396, corresponding to 3.6% of the total particles counted) were
246 identified as polymers, while 9.2% (n=51, i.e. 1% of the total) were pigment particles, such as azure,
247 Hostasol Green G-K, confirming their synthetic nature. The remaining 19% of these particles analyzed
248 could not be identified, probably because of their vivid color or their poor purity. It has already been noted
249 that detecting the chemical composition of the polymer in microplastics can be difficult if the microplastics
250 contain copolymers and additives (Wang et al., 2020). There were 18 polymers, including rubbers, (**Figure**
251 **3, b**) identified in the particles. According to GESAMP (2015) polymers such as PP, PE, PMMA, PVC, PS and
252 PET are often used in consumer goods including packaging, bottles, coatings, films and building materials.
253 No significant differences in the distribution of polymers were observed between sites in this study
254 (Mann-Whitney p-value >0.05).

255 Overall, PA accounted for 24% of the particles analyzed, followed by PET 16% and PC 10%. PLA, which is
256 now used in many consumer goods, was the only bioplastic detected, accounting for 7.5% and 5% at S1
257 and S2 respectively (**Figure 3, b**).

258 The diversity of polymers at Site S2 was greater than at Site 1 as four additional polymers were identified:
259 rubber (0.7%), PMMA (0.7%) PVC (0.4%), and PAEK (0.3%), while PAN (0.7%) was only detected at Site 1.
260 This variation could be explained by the different types of land use and trends in anthropogenic activities.
261 Various domestic, industrial and agricultural uses of plastics contribute to different polymer emissions
262 into the environment, including the release of polyester, and acrylic fibers from textile manufacturing
263 (Napper et al.,2016). The high percentages of PA and PET could also be linked to the discharge of fibers
264 coming from untreated domestic wastewater generally discharged directly into rivers. Su et al. (2020)
265 reported PA to be the main type of microplastic in the rivers and harbors of Melbourne, Australia. These
266 PA and PET fibers are the main constituents of synthetic garments, clothing, tires and mechanical
267 equipment parts (Azziz et al., 2019; Zhang et al., 2019b). This assumption is consistent with our results
268 considering the high proportions of these fibers among the particles.

269

270 2. The impact of hydrological and meteorological conditions on the variation of microplastics

271 a) Abundance of particles according to meteorological conditions

272 The abundance of particles ranged between 0.57 and 28.91 particles/L and between 1.3 and 22.87
273 particles/L during the dry and wet seasons respectively (**Figure 4,a**). No significant differences were

274 highlighted concerning MP abundance between the dry and wet seasons at each site ($p > 0.05$). However,
275 different trends were observed: Site 1 exhibited a significant positive correlation between total MP
276 abundance and daily precipitation (Spearman's rank correlation coefficient, $p < 0.05$), whereas Site 2
277 showed a significant negative correlation between total MP abundance, cumulative precipitation and river
278 flow rate (Spearman's rank correlation coefficient, $p < 0.05$) (**Table 1**). Both sites showed a significant
279 positive correlation between flow and cumulative precipitation. Overall, regarding rainfall events, this
280 study showed lower concentrations compared to a study carried out in Sucy-en-Brie, Paris, across four
281 rainfall events, where MP concentration ranged between 3 and 129 items/L (Treilles et al., 2021a).

282 On the one hand, Site S1 showed a high abundance of particles during the wet season compared to the
283 dry season (**Figure 4, a**), where the flow rate was very low (**Supplementary data, Table 1**). This flow rate
284 of $0.025 \text{ m}^3/\text{s}$ could explain the low influx of particles into the Ibrahim River (S1). The hydrological
285 characteristics of this site could be representative of many of the rivers in Lebanon as they are in actual
286 fact mostly slow-moving streams (Hourri et al., 2007). The highest abundance of particles (4.59 particles/L)
287 was obtained during the first wet season sampling campaign. This can probably be linked to an influx of
288 particles by rain, increasing stormwater runoff and the flow of particles from land to water.

289 Several studies highlighted the occurrence and distribution of MPs in surface waters, influenced by
290 rainfall, with the highest abundance seen during significant precipitation events (Piñon-Colin et al., 2020;
291 Rowley et al., 2020). For example, the highest concentrations of particles in the largest urban lake in China
292 ranged between 7.4 and 29.6 items/L and were directly related to rainfall (Xia et al., 2020). Hitchcock et
293 al. (2020) also observed the strong influence of rainfall, with concentrations rising from 0.4 to 17.38
294 particles/L before and after rainfall. Another study highlighted the contribution of stormwater runoff to
295 an annual load of particles six times greater than the effluent from wastewater treatment plants (Shruti
296 et al. 2021). The impact of stormwater depends on the type of sewer system, whether combined or
297 separate, and on the climatological conditions, as stormwater peak flows can reach high values depending
298 on hydro-meteorological conditions (Treilles et al., 2021b).

299 On the other hand, the Site S2 showed an opposite trend compared to S1 during rainfall events, where a
300 higher abundance of particles was observed during the dry season compared to the wet season, (**Figure**
301 **4, a**). High concentrations during dry seasons could be related to low river flow rates of about $0.16 \text{ m}^3/\text{s}$,
302 (**Supplementary data, Table 1**) as plastic accumulation can be enhanced with low flow rates (Ding et al.,
303 2019). Indeed, other studies have also recorded a high abundance of particles during dry periods and at
304 low flow rates (Bauer-Civiello et al., 2019; de Carvalho et al., 2021; Han et al., 2020; Wang et al., 2021a; Li
305 et al., 2023; Huang et al. 2023). Moreover, a lower surface-water area ratio could occur during the dry

306 season where a river's water volume could be affected, which results in a higher concentration of MPs in
307 the surface water (Wicaksono et al., 2021).

308 In addition, the variation of MP abundance was studied in relation to cumulative precipitation during
309 rainfall episodes (**Figure 4, b**). At the beginning of the wet season, both Sites S1 and S2 showed high
310 abundances of particles. This could be related to the rain washing away any plastics deposited during the
311 dry season and carrying them into the river. For example, a 43-fold increase in the concentration of
312 microplastics (0.4 to 17.383 particles/L) was reported in the Australian Cooks River approximately 12 hrs
313 after it started raining (Hitchcock 2020). Another hypothesis could explain this phenomenon by the
314 vertical exchange occurring between the riverbed, more specifically the active sediment layer, and the
315 watercourses (Ockelford et al. 2020, Kumar et al. 2021). However, for the first sampling campaign during
316 the rainy season at both sites, a higher water level compared to the dry season was clearly observed.
317 Similarly, increased river flow due to rainfall can mobilize sediments, enhancing soil erosion, and this was
318 reported at S1 by Korfali and Davies, (2000). Based on our on-site observations during sampling, high
319 water turbidity was observed at both sites. Hence, following sediment transport, plastics can be
320 resuspended and transported in water (Zhang et al. 2016). As stated by Xia et al. (2021), during
321 resuspension, smaller microplastics (50 μm - 500 μm) may have been displaced from the riverbed into the
322 overlying water.

323 Thereafter, our results showed a progressive decrease in particle abundance as cumulative precipitation
324 (D-10) increased at both sites (**Figure 4, b**). This decrease could be explained by the dilution effect due to
325 rain events, already observed in other studies (Boni et al., 2021; Qian et al., 2023; Wang et al., 2021a).
326 Moreover, the higher flow rate values measured during the last two wet-season campaigns
327 (**Supplementary data, Table 1**) support this hypothesis of a plastic pollution dilution effect. Indeed, the
328 higher volume of water caused by daily and cumulative precipitation could dilute the concentration of
329 microplastics observed.

330 River morphology may also influence the distribution and accumulation of plastic pollution in water
331 ecosystems. However, few studies have described the different convergences of microplastics in river
332 systems due to sites, or channel morphology and topography, e.g. the depth and size of the waterway
333 channel (Crew et al. 2020, Kumar et al. 2021). In this study, two different profiles of rivers stand out
334 concerning their morphology and their sewerage systems which could explain the differences in MP
335 abundance observed.

336 Furthermore, when comparing both sites during the dry season, an opposite trend was observed. The
337 abundance of microplastics at Site S1 progressively decreased, from 1.24 particles/L at the start of the dry
338 season to 0.57 particles/L at the end of the season. In contrast, Site S2 exhibits a progressive increase of
339 MP abundance from 15.54 particles per liter to 28.91 particles per liter over the same period (**Figure 4,**
340 **b**).

341 On the one hand, the Antelias River (S2) is a channeled river (1.8 kilometers long and 6.28 meters wide)
342 with a relatively simple surface profile, featuring infrequent meandering and a fairly free-flowing stream.
343 This channel structure limits the diversity of MP sources, i.e. the interaction between soil and water.
344 Despite the presence of a sewage system to which most buildings are connected, S2 continuously receives
345 untreated sewage from the surrounding area, increasing the risk of the accumulation of MPs (Nel et al.
346 2018). The progressive increase in microplastic abundance at S2 during the dry season can be attributed
347 to the continuous influx of non-treated wastewater into this stream. A dilution of the microplastic
348 accumulation occurred on the first day of precipitation and led to a decrease in MP abundance. Moreover,
349 it is likely that the river morphology limited the resuspension of plastics from the sediment, which also
350 contributes to observations of lower concentrations during these periods of increased runoff and
351 precipitation.

352 On the other hand, the Ibrahim River (S1) is a complex river system (28 kilometers long and 21.79 meters
353 wide) with a free-flowing stream with a more twisted profile and larger dimensions. It is abundantly
354 equipped with septic tanks, suggesting that the water is less affected by direct plastic pollution from the
355 surrounding environment. The abundance seen during the dry season gradually decreases. The highest
356 abundance observed on the first day of precipitation could be linked to septic tank leaching, transporting
357 microplastics into the river.

358 b) Morphological and polymeric variations of MPs by season

359 Fibers and films are significantly more abundant during the dry season at S2 compared to the dry season
360 at S1 (Kruskal-Wallis, $p < 0.05$) (**Figure 5, a**). A higher proportion of fragments tended to be observed
361 during the wet season at both sites ($12\%_{\text{dry}} < 21\%_{\text{wet}}$). Fragments showed significantly higher abundance
362 during the wet season exclusively at S1 (Kruskal-Wallis, $p < 0.05$). A significant positive correlation was
363 found between microbeads and cumulative precipitation and flow at S1. In contrast, a significant negative
364 correlation was seen between films and cumulative precipitation and flow, as well as a similar significant
365 negative correlation between fibers and cumulative precipitation and flow at S2 (**Table 1**). A similar shape
366 distribution was observed in runoff waters from Tijuana, Mexico (Piñon-Colin et al., 2020) and Gündoğdu

367 et al. (2018) reported a significant increase in film after the flood. Particles smaller than 1 mm always
 368 represented the main size class, regardless of meteorological conditions, **(Supplementary data, Figure 2)**.
 369 This could be explained by the effect of the precipitation that can carry the smallest particles trapped into
 370 the surface layer of the soil, transferring them into stormwater (Shruti et al., 2021; Zhang et al., 2017).
 371 More polymers were identified during the wet season than during the dry season. At each site, the
 372 proportion of PET tended to double during the dry season 29.4% (S1) and 25.6% (S2) compared to the wet
 373 season 11.8% (S2) and 15.6% (S1) **(Figure 5, b)**. Among the polymers identified in the samples from S1,
 374 PAN and PP-PE were only identified during the wet season, while PUR was detected only during the dry
 375 season. At Site S2, PMMA, PE, PVC and rubber were only identified during wet season whereas PAEK and
 376 PUR were only present during dry season. The rainfall at Site 2 may have favored the transportation of
 377 these higher-density polymers. These observations could highlight the role of meteorological
 378 characteristics, as either rain or stormwater runoff can be a vector for transporting new polymers to the
 379 receiving site (Hitchcock 2020).

380 c) Flow of microplastics into the Mediterranean Sea

381 Daily water flows at each site were provided by the Litani River Authority **(Supplementary data, Table 1)**
 382 and used to calculate the flux of microplastics per day and per season.

383 The following formula was used to calculate the flux of MPs per second at each site:

$$384 \quad (1) \quad Flux_{particles/s} = Abundance_{\frac{particles}{m^3}} \times Daily\ Flow_{\frac{m^3}{s}}$$

385 The following formula was used to calculate the flux of MPs per day at each site:

$$386 \quad (2) \quad Flux_{particles/day} = Abundance_{\frac{particles}{m^3}} \times Daily\ Flow_{\frac{m^3}{s}} \times 60 \times 60 \times 24$$

387 The estimated flux of MPs during the dry period at S2 shows higher values ranging between 2.15×10^5
 388 and 4×10^5 particles/day compared to S1 values ranging between 1.23×10^3 and 2.68×10^3 particles/day.
 389 This is because S2 has a low daily flow rate and a higher abundance of MPs during the dry season. This
 390 result suggests that during dry periods, the most urbanized sites contribute the most to the sea pollution.

391 Moreover, the estimated flow of MPs during the rainy season at S1 has higher values (between 1.07×10^5
 392 and 2.29×10^6 particles/day) than at S2 (between 1.64×10^5 and 3.56×10^5 particles/day). These results
 393 highlight the effect of hydrological characteristics, with the river having a higher flow rate during the wet
 394 season, and hence contributing more to plastic pollution in the sea.

395 In this study, we have emphasized the fact that untreated wastewater in Lebanon is a major source of
396 microplastics in rivers that enter the Mediterranean Sea largely during the dry season. Hence, hydrological
397 and meteorological conditions enhance the flux of MPs from rivers to the sea in regions with lower levels
398 of anthropogenic activities. Anthropogenic activities are therefore considered to be a direct and indirect
399 source of primary plastic pollution, since plastics are used intentionally on a daily basis (Napper and
400 Thompson 2016, de Carvalho et al. 2021).

401

402 IV. Conclusion

403 To our knowledge, this study is the first that has described the contamination of MPs in Lebanese rivers
404 along the Mediterranean Sea. Microplastics were found in all surface water samples and our results
405 revealed that plastic pollution is significantly positively influenced by the degree of urbanization around
406 these rivers both during the dry and wet seasons. However, the effect of meteorological conditions on
407 the abundance of MPs appeared contradictory, depending on the site. In the small river at S2, where
408 anthropogenic influence is greater, we observed high concentrations of microplastics during periods of
409 low flow. In contrast, the larger river (S1), characterized by its larger size and higher flow rate volume,
410 tended to show higher pollution levels during the wet season. These differences are consistent with
411 differences in anthropogenic activities, hydrologic conditions, and sewage management activities. Our
412 results also showed a higher abundance during the first rain event at both sites, followed by a gradual
413 decrease that tends to follow cumulative rainfall, suggesting a pollution dilution effect. In all samples, the
414 particles were mostly in the form of fibers smaller than 1 mm. These particles were mainly comprised of
415 PA and PET but some differences in polymer composition were identified between the seasons at each
416 site.

417 The flows of the MP contamination calculated has shown the importance of small river discharges into
418 the Mediterranean Sea and has highlighted the influence of hydrological and meteorological conditions
419 on their contribution to coastal pollution.

420 V. Credit Roles

421 **Rosa Sawan:** Conceptualization, Methodology, Investigation, Formal analysis, Data curation, Writing -
422 original draft; **Périne Doyen:** Conceptualization, Supervision, Visualization, Writing - review & editing;
423 **Florence Viudes:** Investigation; **Rachid Amara:** Conceptualization, Methodology, Supervision, Funding
424 acquisition, Visualization, Writing - review & editing; **Céline Mahfouz:** Conceptualization, Methodology,
425 Supervision, Funding acquisition, Visualization, Writing - review & editing.

426 **VI. Acknowledgments**

427 The authors would like to thank M. Elie Tarek, for its help during field work, and the National Center for
428 remote sensing (CNRS-Liban) for providing the land cover data. We would also like to thank the
429 Laboratoire de Sécurité des Aliments of Anses for their help with μ FTIR analyses.

430

431 **VII. Funding**

432 The authors are very grateful to ULCO (University of Littoral Côte d'Opale) and National Center of Marine
433 Sciences of Lebanon (CNRS-Lebanon) for financially supporting Rosa Sawan's PhD scholarship. This work
434 has benefited from the grant "ANR-21-EXES-00 11" as part of the IFSEA graduate school, which originates
435 from National Research Agency under the Investments for the Future program. This work has been also
436 financially by the SFR Campus de la Mer and by the European Union (ERDF), the French Government, the
437 Région Hauts-de-France and IFREMER, in the framework of the project CPER IDEAL 2021-2027.

438 VII. References

- 439 Abidli S, Antunes JC, Ferreira JL, Lahbib Y, Sobral P, Trigui El Menif N. 2018. Microplastics in
440 sediments from the littoral zone of the north Tunisian coast (Mediterranean Sea). *Estuarine, Coastal
441 and Shelf Science* 205: 1–9.
- 442 Adomat Y, Grischek T. 2021. Sampling and processing methods of microplastics in river sediments - A
443 review. *Science of The Total Environment* 758: 143691.
- 444 Alomar C, Estarellas F, Deudero S. 2016. Microplastics in the Mediterranean Sea: Deposition in coastal
445 shallow sediments, spatial variation and preferential grain size. *Marine Environmental Research*
446 115: 1–10.
- 447 Assaker A. 2016. Hydrologie et biogéochimie du bassin versant du fleuve Ibrahim : un observatoire du
448 fonctionnement de la zone critique au Liban (Hydrology and biogeochemistry of the Ibrahim River
449 Basin: an observatory of the critical zone functioning in Lebanon). 1–278.
- 450 Auta HS, Emenike CU, Fauziah SH. 2017. Distribution and importance of microplastics in the marine
451 environment: A review of the sources, fate, effects, and potential solutions. *Environment
452 international* 102: 165–176.
- 453 Azziz Assoumani, Marie-Pierre Strub, Sophie Lardy-Fontan, Enrica Alasonati FG. 2019. Microplastiques
454 dans les eaux de surface continentales. Rapport AQUAREF 63.
- 455 Balthazar-Silva D, Ft M, Turra A, Moreira F, Camargo R, Oliveira A, Barbosa L, Gorman D. 2020.
456 Rainfall and Tidal Cycle Regulate Seasonal Inputs of Microplastic Pellets to Sandy Beaches.
457 *Frontiers in Environmental Science* 8: 1–8.
- 458 Bauer-Civiello A, Critchell K, Hoogenboom M, Hamann M. 2019. Input of plastic debris in an urban
459 tropical river system. *Marine Pollution Bulletin* 144: 235–242.
- 460 Boni W, Arbuckle-Keil G, Fahrenfeld NL. 2021. Inter-storm variation in microplastic concentration and
461 polymer type at stormwater outfalls and a bioretention basin. *The Science of the total environment*
462 151104.
- 463 Campanale C, Stock F, Massarelli C, Kochleus C, Bagnuolo G, Reifferscheid G, Uricchio VF. 2020.
464 Microplastics and their possible sources: The example of Ofanto river in southeast Italy.
465 *Environmental Pollution* 258: 113284.
- 466 de Carvalho AR, Garcia F, Riem-Galliano L, Tudesque L, Albignac M, ter Halle A, Cucherousset J. 2021.
467 Urbanization and hydrological conditions drive the spatial and temporal variability of microplastic

468 pollution in the Garonne River. *Science of The Total Environment* 769: 144479.

469 Celine M, Sharif J, Maria K, El Rahman HA, Myriam L, Myriam G, Anthony O, Rachid A, Milad F.
 470 2023. First assessment of microplastics in offshore sediments along the Lebanese coast, South-
 471 Eastern Mediterranean. *Marine Pollution Bulletin* 186: 114422.

472 Chen J, Deng Y, Chen Y, Peng X, Qin H, Wang T, Zhao C. 2022. Distribution Patterns of Microplastics
 473 Pollution in Urban Fresh Waters: A Case Study of Rivers in Chengdu, China. *International Journal*
 474 *of Environmental Research and Public Health* 19: .

475 Cincinelli A, Martellini T, Guerranti C, Scopetani C, Chelazzi D, Giarrizzo T. 2019. A potpourri of
 476 microplastics in the sea surface and water column of the Mediterranean Sea. *TrAC Trends in*
 477 *Analytical Chemistry* 110: 321–326.

478 Constant M, Ludwig W, Kerhervé P, Sola J, Charrière B, Sanchez-Vidal A, Canals M, Heussner S. 2020.
 479 Microplastic fluxes in a large and a small Mediterranean river catchments: The Têt and the Rhône,
 480 Northwestern Mediterranean Sea. *Science of The Total Environment* 716: 136984.

481 Crew A, Gregory-Eaves I, Ricciardi A. 2020. Distribution, abundance, and diversity of microplastics in
 482 the upper St. Lawrence River. *Environmental Pollution* 260: 113994.

483 Daou C, Salloum M, Mouneimne AH, Legube B, Ouaini N. 2013. Multidimensionnal analysis of two
 484 Lebanese surface water quality: Ibrahim and el-Kalb rivers. *Journal of Applied Sciences Research* 9:
 485 2777–2787.

486 Demir A. 2022. Riverine Microplastic Loading to Mersin Bay, Turkey on the North-eastern
 487 Mediterranean. *Turkish Journal of Fisheries and Aquatic Sciences* 21: .

488 Ding L, Mao R fan, Guo X, Yang X, Zhang Q, Yang C. 2019. Microplastics in surface waters and
 489 sediments of the Wei River, in the northwest of China. *Science of The Total Environment* 667: 427–
 490 434.

491 Dris R, Gasperi J, Rocher V, Mohamed S, Tassin B. 2015a. Microplastiques: sources et milieux
 492 récepteurs dans un contexte urbain. .

493 Dris R, Gasperi J, Rocher V, Saad M, Renault N, Tassin B. 2015b. Microplastic contamination in an
 494 urban area: a case study in Greater Paris. *Environmental Chemistry* 12: 592–599.

495 Eerkes-Medrano, Thompson, Aldridge. 2015. Microplastics in freshwater systems: a review of the
 496 emerging threats, identification of knowledge gaps and prioritisation of research needs. *Water*

497 research 75: 63–82.

498 van Emmerik T, Kieu-Le T-C, Loozen M, van Oeveren K, Strady E, Bui X-T, Egger M, Gasperi J,
499 Lebreton L, Nguyen P-D, Schwarz A, Slat B, Tassin B. 2018. A Methodology to Characterize
500 Riverine Macroplastic Emission Into the Ocean . *Frontiers in Marine Science* 5: .

501 Fagiano V, Compa M, Alomar C, Rios-Fuster B, Morató M, Capó X, Deudero S. 2023. Breaking the
502 paradigm: Marine sediments hold two-fold microplastics than sea surface waters and are dominated
503 by fibers. *Science of the Total Environment* 858: .

504 Falah A, Nenov V, Yemendzhiev H. 2021. Inland Water Quality Study of Mount Lebanon Range in the
505 Region of Middle East. .

506 GESAMP. 2015. Sources, fate and effects of microplastics in the marine environment: A global
507 assessment. .

508 Gündoğdu S, Çevik C, Ayat B, Aydoğan B, Karaca S. 2018. How microplastics quantities increase with
509 flood events? An example from Mersin Bay NE Levantine coast of Turkey. *Environmental
510 pollution (Barking, Essex : 1987)* 239: 342–350.

511 Hakim B. et Karkabi S. 1988. Colorations du gouffre de Faouar Dara et de la grotte des Kassarat (Caza du
512 Metn Nord-Liban) [Colorations of sinkhole of Faouar Dara and the grotto of Kassarat (Metn
513 District North Lebanon)] 1965- 1967-1969. *AL OUAT'OUATE Revue Libanaise de Spéléologie et
514 de Karstologie.* 18–31.

515 Han M, Niu X, Tang M, Zhang B-T, Wang G, Yue W, Kong X, Zhu J. 2020. Distribution of
516 microplastics in surface water of the lower Yellow River near estuary. *The Science of the total
517 environment* 707: 135601.

518 Hanna N, Lartiges B, Kazpard V, Maatouk E, Amacha N, Sassine S, El Samrani A. 2018.
519 Hydrogeochemical Processes in a Small Eastern Mediterranean Karst Watershed (Nahr Ibrahim,
520 Lebanon). *Aquatic Geochemistry* 24: 325–344.

521 He D, Chen X, Zhao W, Zhu Z, Qi X, Zhou L, Chen W, Wan C, Li D, Zou X, Wu N. 2021. Microplastics
522 contamination in the surface water of the Yangtze River from upstream to estuary based on
523 different sampling methods. *Environmental research* 196: 110908.

524 Hitchcock JN. 2020. Storm events as key moments of microplastic contamination in aquatic ecosystems.
525 *The Science of the total environment* 734: 139436.

- 526 Hitchcock JN, Mitrovic SM. 2019. Microplastic pollution in estuaries across a gradient of human impact.
527 Environmental Pollution 247: 457–466.
- 528 Horton, Walton, Spurgeon, Lahive E, Svendsen C. 2017. Microplastics in freshwater and terrestrial
529 environments: Evaluating the current understanding to identify the knowledge gaps and future
530 research priorities. Science of The Total Environment 586: 127–141.
- 531 Hourani A, Galev E, Chahine M. 2019. Decompression versus compression Fouar Antelias: A gushing
532 water river. AGROFOR 3: .
- 533 Houri A, El Jeblawi SW. 2007. Water quality assessment of Lebanese coastal rivers during dry season
534 and pollution load into the Mediterranean Sea. Journal of water and health 5: 615–623.
- 535 Huang C-W, Li Y-L, Lin C, Bui X-T, Vo T-D-H, Ngo HH. 2023. Seasonal influence on pollution index
536 and risk of multiple compositions of microplastics in an urban river. Science of The Total
537 Environment 859: 160021.
- 538 Hurley R, Woodward J, Rothwell JJ. 2018. Microplastic contamination of river beds significantly reduced
539 by catchment-wide flooding. Nature Geoscience 11: 251–257.
- 540 Jemaa S, Mahfouz C, Kazour M, Lteif M, El A, Hassoun AER, Ghsoub M, Amara R, Khalaf G, Fakhri
541 M. 2021. Floating Marine Litter in Eastern Mediterranean From Macro to Microplastics: The
542 Lebanese Coastal Area as a Case Study. Frontiers in Environmental Science 9: .
- 543 Kazour M, Jemaa S, Issa C, Khalaf G, Amara R. 2019. Microplastics pollution along the Lebanese coast
544 (Eastern Mediterranean Basin): Occurrence in surface water, sediments and biota samples. Science
545 of The Total Environment 696: 133933.
- 546 Khatib J, Baydoun S, Elkordi A. 2018. Water Pollution and Urbanisation Trends in Lebanon: Litani River
547 Basin Case Study: Science and Management. Urban Pollution: Science and Management p. 397–
548 415.
- 549 Korfali SI, Davies BE. 2000. Total and Extractable Trace Elements in Lebanese River Sediments: Dry
550 Season Data. Environmental Geochemistry and Health 22: 265–273.
- 551 Kumar R, Sharma P, Verma A, Jha PK, Singh P, Gupta P, Chandra R, Prasad PVV. 2021. Effect of
552 Physical Characteristics and Hydrodynamic Conditions on Transport and Deposition of
553 Microplastics in Riverine Ecosystem. Water 13: 2710.
- 554 Kundu MN, Komakech HC, Lugomela G. 2022. Analysis of Macro- and Microplastics in Riverine,

555 Riverbanks, and Irrigated Farms in Arusha, Tanzania. *Archives of Environmental Contamination*
556 *and Toxicology* 82: 142–157.

557 Kunz A, Schneider F, Anthony N, Lin H-T. 2023. Microplastics in rivers along an urban-rural gradient in
558 an urban agglomeration: Correlation with land use, potential sources and pathways. *Environmental*
559 *Pollution* 321: 121096.

560 Li J, Liu H, Paul Chen J. 2018. Microplastics in freshwater systems: A review on occurrence,
561 environmental effects, and methods for microplastics detection. *Water Research* 137: 362–374.

562 Li T, Liu K, Tang R, Liang J-R, Mai L, Zeng EY. 2023. Environmental fate of microplastics in an urban
563 river: Spatial distribution and seasonal variation. *Environmental Pollution* 322: 121227.

564 Li Y, Lu Z, Zheng H, Wang J, Chen C. 2020. Microplastics in surface water and sediments of Chongming
565 Island in the Yangtze Estuary, China. *Environmental Sciences Europe* 32: 15.

566 Lopes C, Raimundo J, Caetano M, Garrido S. 2020. Microplastic ingestion and diet composition of
567 planktivorous fish. *Limnology and Oceanography Letters* 5: .

568 Luo W, Su L, Craig NJ, Du F, Wu C, Shi H. 2019. Comparison of microplastic pollution in different
569 water bodies from urban creeks to coastal waters. *Environmental Pollution* 246: 174–182.

570 Maatouk E. 2014. Caractérisation des eaux usées au Liban : impact sur le fonctionnement des stations
571 d'épuration.

572 Maatouk E, Samrani A El, Sawan R, Salameh RBG, Kazpard V, Matar Z. 2022. Influence of Diverse
573 Urban Pressures on Water Characteristics in a Small Eastern Mediterranean Watershed. *Aquatic*
574 *Geochemistry* 28: 111–133.

575 El Najjar P, Kassouf A, Probst A, Probst JL, Ouaini N, Daou C, El Azzi D. 2019. High-frequency
576 monitoring of surface water quality at the outlet of the Ibrahim River (Lebanon): A multivariate
577 assessment. *Ecological Indicators* 104: 13–23.

578 El Najjar P, Pfaffl M, Ouaini N, Abdel Nour A, El Azzi D. 2020. Water and sediment microbiota
579 diversity in response to temporal variation at the outlet of the Ibrahim River (Lebanon).
580 *Environmental Monitoring and Assessment* 192: 201.

581 Napper IE, Thompson RC. 2016. Release of synthetic microplastic plastic fibres from domestic washing
582 machines: Effects of fabric type and washing conditions. *Marine Pollution Bulletin* 112: 39–45.

583 Nehme C, Jaillet S, Delannoy J-J, Gerard J, Hellstrom J. 2013. Sedimentary study and U-TH datations

584 contribution to the morphodynamic reconstitution of the junction chamber (Kassarat cave, Nabay,
585 Lebanon): A geomorphological approach for palaeohydrological records analysis.

586 Nel HA, Dalu T, Wasserman RJ. 2018. Sinks and sources: Assessing microplastic abundance in river
587 sediment and deposit feeders in an Austral temperate urban river system. *Science of The Total*
588 *Environment* 612: 950–956.

589 Ockelford A, Cundy A, Ebdon JE. 2020. Storm Response of Fluvial Sedimentary Microplastics.
590 *Scientific Reports* 10: 1865.

591 Panti C, Giannetti M, Bains M, Rubegni F, Minutoli R, Fossi MC. 2015. Occurrence, relative abundance
592 and spatial distribution of microplastics and zooplankton NW of Sardinia in the Pelagos Sanctuary
593 Protected Area, Mediterranean Sea. *Environmental Chemistry* 12: 618–626.

594 Park H, Park B. 2021. Review of Microplastic Distribution, Toxicity, Analysis Methods, and Removal
595 Technologies. *Water* 13: .

596 Piñon-Colin T de J, Rodriguez-Jimenez R, Rogel-Hernandez E, Alvarez-Andrade A, Wakida FT. 2020.
597 Microplastics in stormwater runoff in a semiarid region, Tijuana, Mexico. *Science of The Total*
598 *Environment* 704: 135411.

599 Qian Y, Shang Y, Zheng Y, Jia Y, Wang F. 2023. Temporal and spatial variation of microplastics in
600 Baotou section of Yellow River, China. *Journal of Environmental Management* 338: 117803.

601 Rayadin Y, Syamsudin J, Ayatussurur M, Qomari N, Pradesta H, Priahutama, A& Putri RO. 2016.
602 Scholar (8). Convergence in the information industries. *Telecommunications, broadcasting and data*
603 *processing* 1981-1996 3: 45.

604 Rios Mendoza LM, Balcer M. 2019. Microplastics in freshwater environments: A review of quantification
605 assessment. *TrAC Trends in Analytical Chemistry* 113: 402–408.

606 Rodrigues MO, Abrantes N, Gonçalves FJM, Nogueira H, Marques JC, Gonçalves AMM. 2018. Spatial
607 and temporal distribution of microplastics in water and sediments of a freshwater system (Antuã
608 River, Portugal). *Science of The Total Environment* 633: 1549–1559.

609 Rowley K, Cucknell A, Smith B, Clark P, Morrith D. 2020. London’s river of plastic: High levels of
610 microplastics in the Thames water column. *Science of The Total Environment* 740: 140018.

611 Ruiz-Orejón LF, Sardá R, Ramis-Pujol J. 2016. Floating plastic debris in the Central and Western
612 Mediterranean Sea. *Marine Environmental Research* 120: 136–144.

613 Sarijan S, Azman S, Said MIM, Jamal MH. 2021. Microplastics in freshwater ecosystems: a recent
614 review of occurrence, analysis, potential impacts, and research needs. *Environmental science and*
615 *pollution research international* 28: 1341–1356.

616 Shruti VC, Pérez-Guevara F, Elizalde-Martínez I, Kutralam-Muniasamy G. 2021. Current trends and
617 analytical methods for evaluation of microplastics in stormwater. *Trends in Environmental*
618 *Analytical Chemistry* 30: e00123.

619 Simon-Sánchez L, Grelaud M, Garcia-Orellana J, Ziveri P. 2019. River Deltas as hotspots of microplastic
620 accumulation: The case study of the Ebro River (NW Mediterranean). *The Science of the total*
621 *environment* 687: 1186–1196.

622 Slim K, Saad Z, Kazpard V, El Samarani A. 2005. Chemical and Isotopic Composition of Rain water in
623 Coastal and Highland Regions in Lebanon. *Journal of Environmental Hydrology* 1–11.

624 Stovall JK, Bratton SP. 2022. Microplastic Pollution in Surface Waters of Urban Watersheds in Central
625 Texas, United States: A Comparison of Sites With and Without Treated Wastewater Effluent .
626 *Frontiers in Analytical Science* 2: .

627 Su L, Sharp SM, Pettigrove VJ, Craig NJ, Nan B, Du F, Shi H. 2020. Superimposed microplastic
628 pollution in a coastal metropolis. *Water Research* 168: 115140.

629 Talbot R, Chang H. 2022. Microplastics in freshwater: A global review of factors affecting spatial and
630 temporal variations. *Environmental Pollution* 292: 118393.

631 The World Bank TWBG. 2021. [Internet]. Available from:
632 <https://climateknowledgeportal.worldbank.org/country/lebanon/climate-data-historical>

633 Treilles R, Gasperi J, Gallard A, Saad M, Dris R, Partibane C, Breton J, Tassin B. 2021a. Microplastics
634 and microfibers in urban runoff from a suburban catchment of Greater Paris. *Environmental*
635 *Pollution* 287: 117352.

636 Treilles R, Gasperi J, Tramoy R, Dris R, Gallard A, Partibane C, Tassin B. 2021b. Microplastic and
637 microfiber fluxes in the Seine River: Flood events versus dry periods. *Science of The Total*
638 *Environment* 805: 150123.

639 Wang C, Xing R, Sun M, Ling W, Shi W, Cui S, An L. 2020. Microplastics profile in a typical urban
640 river in Beijing. *Science of The Total Environment* 743: 140708.

641 Wang G, Lu J, Li W, Ning J, Zhou L, Tong Y, Liu Z, Zhou H, Xiayihazi N. 2021a. Seasonal variation

642 and risk assessment of microplastics in surface water of the Manas River Basin, China.
643 *Ecotoxicology and environmental safety* 208: 111477.

644 Wang T, Wang J, Lei Q, Zhao Y, Wang L, Wang X, Zhang W. 2021b. Microplastic pollution in
645 sophisticated urban river systems: Combined influence of land-use types and physicochemical
646 characteristics. *Environmental Pollution* 287: 117604.

647 Wang W, Ndungu AW, Li Z, Wang J. 2017. Microplastics pollution in inland freshwaters of China: A
648 case study in urban surface waters of Wuhan, China. *The Science of the total environment* 575:
649 1369–1374.

650 Wicaksono EA, Werorilangi S, Galloway TS, Tahir A. 2021. Distribution and Seasonal Variation of
651 Microplastics in Tallo River, Makassar, Eastern Indonesia. *Toxics* 9: .

652 Wong G, Löwemark L, Kunz A. 2020a. Microplastic pollution of the Tamsui River and its tributaries in
653 northern Taiwan: Spatial heterogeneity and correlation with precipitation. *Environmental Pollution*
654 260: 113935.

655 Wong JKH, Lee KK, Tang KHD, Yap P-S. 2020b. Microplastics in the freshwater and terrestrial
656 environments: Prevalence, fates, impacts and sustainable solutions. *Science of The Total*
657 *Environment* 719: 137512.

658 Wu P, Tang Y, Dang M, Wang S, Jin H, Liu Y, Jing H, Zheng C, Yi S, Cai Z. 2019. Spatial-Temporal
659 Distribution of Microplastics in Surface Water and Sediments of Maozhou River within
660 Guangdong-Hong Kong-Macao Greater Bay Area. *Science of The Total Environment* 717: 135187.

661 Xia F, Yao Q, Zhang J, Wang D. 2021. Effects of seasonal variation and resuspension on microplastics in
662 river sediments. *Environmental Pollution* 286: 117403.

663 Xia W, Rao Q, Deng X, Chen J, Xie P. 2020. Rainfall is a significant environmental factor of microplastic
664 pollution in inland waters. *The Science of the total environment* 732: 139065.

665 Zhang J, Zhang C, Deng Y, Wang R, Ma E, Wang J, Bai J, Wu J, Zhou Y. 2019a. Microplastics in the
666 surface water of small-scale estuaries in Shanghai. *Marine pollution bulletin* 149: 110569.

667 Zhang K, Hamidian AH, Tubić A, Zhang Y, Fang JKH, Wu C, Lam PKS. 2021. Understanding plastic
668 degradation and microplastic formation in the environment: A review. *Environmental Pollution* 274:
669 116554.

670 Zhang K, Su J, Xiong X, Wu X, Wu C, Liu J. 2016. Microplastic pollution of lakeshore sediments from

- 671 remote lakes in Tibet plateau, China. *Environmental Pollution* 219: 450–455.
- 672 Zhang K, Xiong X, Hu H, Wu C, Bi Y, Wu Y, Zhou B, Lam PKS, Liu J. 2017. Occurrence and
673 Characteristics of Microplastic Pollution in Xiangxi Bay of Three Gorges Reservoir, China.
674 *Environmental Science & Technology* 51: 3794–3801.
- 675 Zhang S, Wang J, Liu X, Qu F, Wang X, Wang X, Li Y, Sun Y. 2019b. Microplastics in the environment:
676 A review of analytical methods, distribution, and biological effects. *TrAC Trends in Analytical*
677 *Chemistry* 111: 62–72.
- 678 Zhang Y, Kang S, Allen S, Allen D, Gao T, Sillanpää M. 2020. Atmospheric microplastics: A review on
679 current status and perspectives. *Earth-Science Reviews* 203: 103118.
- 680

681 IX. Table and Figure captions

682 **Table 1** Spearman's rank correlation coefficient matrix between environmental variables and total
683 abundance and types of microplastics at S1 (Ibrahim River) and S2 (Antelias River). p-values < 0.05 are
684 highlighted in bold. The blue color refers to significant positive correlation and the red color refers to
685 significant negative correlation.


686 **Figure 1.** Location of the two areas studied along the Lebanese coast, the Eastern Mediterranean Sea.
687 Land cover usage maps of the Ibrahim River outlet (S1) and the Antelias River outlet (S2), indicating the
688 surface of urban and industrial areas and the natural area.

689 **Figure 2.** Variation of particles abundance and morphological characteristics at each site Ibrahim (S1) and
690 Antelias (S2): (a) Boxplot showing of abundance, where the mean abundance is indicated by an X inside
691 of each box; and the median is indicated by the line inside, while bars are for minimum and maximum
692 abundance, (b) the percentage of particle types (fibers, films, fragments and microbeads), (c) particle size
693 (A, S1), and (B, S2). The size scale in μm was divided into four groups (150-500, 500-1,000, 1,000-3,000 and
694 3,000-5,000). (C) and (D) plot the variability of MP types over the 150 μm – 500 μm size range. (d) color
695 distribution of particles (in %).



696 **Figure 3(a).** Examples of spectra obtained when identifying the type of polymer: (A)PA, (B)PET.
697 (b)Percentage of polymer distribution at both sites: the Ibrahim River (S1) and the Antelias River (S2).
698 Polyamide (PA,nylon), polyethylene terephthalate (PET), polycarbonate (PC), polystyrene (PS, ePS),
699 acrylonitrile butadiene styrene (ABS), polypropylene (PP), polylactic acid (PLA), polyurethane (PUR),
700 polyethylene (PE), copolymer like polypropylene-polyethylene (PP-PE), polymethyl methacrylate (PPMA),
701 polyvinyl chloride (PVC), cellulose acetate (CA), P-aryletherketone (PAEK), polyacrinortile (PAN),
702 polytetrafluoroethylene (PTFE) and rubber.







703 **Figure 4.** Variation of particles abundances at each site Ibrahim (S1) and Antelias (S2) during the dry and
704 wet season. (a) Boxplot showing the variation of the particle concentration, where the mean abundance
705 is indicated by an X inside of each box; bars are for minimum and maximum abundance. (b) Bar charts
706 S1(orange bars), S2(blue bars), during each sampling period as a function of daily (D dashed line) and
707 cumulative precipitation (D-10 dark blue line).



708 **Figure 5.** (a)Variation in the percentage of particle types (fibers, films, fragments and microbeads) at both
709 sites (the Ibrahim River, S1 and the Antelias River, S2) during the wet and dry seasons. (b) Polymer
710 distribution during the wet and dry season at both sites S1 (the Ibrahim River) and S2 (the Antelias River).









Legend




-  Ibrahim Outlet
-  Antelias Outlet
-  Rivers



-  Agriculture
-  Bare Rocks
-  Dense Urban and touristic areas,
-  Diverse Equipment, anthropogenic activities
-  Highway
-  Hill Lake
-  Industrial Areas
-  Low Density Urban Areas
-  Medium Density Urban Areas
-  Mineral Extraction Site
-  Port Areas
-  River
-  Sandy Beaches
-  Lebanon











	Daily Precipitation	Cumulative Precipitation	Flow rate	Total Abundance	Fibers	Films	Fragments	Beads
S1								
Daily Precipitation	1							
Cumulative Precipitation	0.143	1						
Flow rate	0.143	0.000	1					
Total Abundance	0.035	0.129	0.129	1				
Fibers	0.114	0.518	0.519	0.014	1			
Films	0.096	0.742	0.742	0.077	0.007	1		
Fragments	0.114	0.161	0.161	0.003	0.094	0.233	1	
Beads	0.194	0.006	0.006	0.332	0.758	1.000	0.332	1
S2								
Daily Precipitation	1							
Cumulative Precipitation	0.143	1						
Flow rate	0.143	0.000	1					
Total Abundance	0.579	0.039	0.039	1				
Fibers	0.579	0.039	0.039	0.001	1			
Films	0.261	0.007	0.007	0.014	0.052	1		
Fragments	0.240	0.691	0.691	0.180	0.294	0.294	1	
Beads	0.958	0.547	0.547	0.049	0.049	0.317	0.216	1