

# Temporal and Spatial Dynamics of Vibrio Harveyi: An Environmental Parameter Correlated Investigation in a 4-Metre-Deep Aquaculture Tank

Alix da Fonseca Ferreira, Roxane Roquigny, Thierry Grard, Cédric Le Bris

# ▶ To cite this version:

Alix da Fonseca Ferreira, Roxane Roquigny, Thierry Grard, Cédric Le Bris. Temporal and Spatial Dynamics of Vibrio Harveyi: An Environmental Parameter Correlated Investigation in a 4-Metre-Deep Aquaculture Tank. Microorganisms, 2024, 12 (6), pp.1104. 10.2139/ssrn.4777236 . hal-04592601

# HAL Id: hal-04592601 https://ulco.hal.science/hal-04592601

Submitted on 29 May 2024  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

# <sup>1</sup> Temporal and spatial dynamics of Vibrio

- <sup>2</sup> *harveyi*: an environmental parameter
- <sup>3</sup> correlated investigation in a 4-metre-deep

# aquaculture tank

- 5 A. Da Fonseca Ferreira, R. Roquigny, T. Grard and C. Le Bris
- 6 Univ. Littoral Côte d'Opale, UMRt 1158 BioEcoAgro, USC ANSES, INRAe, Univ. Artois, Univ.Lille, Univ. Picardie Jules
- 7 Verne, Univ. Liège, Junia, 62200 Boulogne-sur-Mer, France

# 8 Abstract

9 Nowadays, European sea bass (Dicentrarchus labrax) aquaculture knows a significant expansion. 10 Nevertheless, the aquaculture industry is plagued by several microbial diseases, including vibriosis. The spatial and temporal dynamics of pathogenic Vibrio harveyi were studied on a European seabass farm in 11 12 northern France over seven months in 2022. Cylindrical concrete specimens were suspended in a water column and water was pumped from different depths (top: 0.3 m deep, middle: 2.15 m deep, and bottom: 13 14 4 m deep), providing insights into biofilm and planktonic V. harveyi dynamics. The abundances of V. 15 harveyi, in biofilm and free-living forms, were positively correlated. Water parameters were monitored, 16 revealing seasonal fluctuations in temperature, pH, and salinity, with no significant differences observed 17 across the water column. Fish mortality peaks were recorded in July and August, coinciding with elevated 18 temperatures and emphasizing the impact of environmental factors on aquaculture health. Quantification 19 of V. harveyi by quantitative PCR revealed no significant differences between water top, middle and 20 bottom layers. Vibrio harveyi exhibited seasonality, with peak abundances observed in August, correlating 21 with temperature increases. Principal component analysis identified temperature as a primary driver, but also additional parameters such as salinity and pH. vibriosis outbreaks occurred during the sampling 22 23 period, providing valuable insights on conditions before, during, and after the outbreak. This study 24 underscores the importance of understanding V. harveyi behaviour in aquaculture systems, particularly 25 in the context of global warming, for effective disease management and sustainable practices.

## 26 Keywords

28

27 Aquaculture, Vibrio harveyi, Biofilms, Planktonic, Environment, Spatio-temporal study

# 1. Introduction

29 Given the swift expansion of the global population and the sharp decrease in catches from marine 30 fisheries, the advancement of aquaculture has gained substantial attention as a means to guarantee 31 worldwide food supplies (Ahmed et al., 2019). The European seabass (Dicentrarchus labrax) stands out as one of the most economically significant fish species harvested in Europe. Over recent decades, the 32 33 European seabass aquaculture sector has experienced robust growth. Overall production has reached 34 80,786 tonnes, amounting to a value of € 470 million in 2020 (EUMOFA, 2022). However, like many other 35 species, D. labrax is susceptible to various bacterial diseases, including pasteurellosis (Photobacterium damselae subsp. piscicida), photobacteriosis (Photobacterium damselae subsp. damselae), 36 37 tenacibaculosis (Tenacibaculum spp.), and vibriosis (Vibrio spp.). These diseases pose significant threats to the seabass aquaculture sector (Muniesa et al., 2020). They can not only cause mass mortality events, 38 39 but also have substantial economic consequences (ANSES, 2019; Novriadi, 2016).

Throughout the whole production chain, vibriosis stands out as the most significant threat to seabass. It
 is characterised by various symptoms in seabass, including anorexia, erratic swimming behaviour,
 exophthalmia, darkening of body colour, pale gills, haemorrhage, ulcers, and skin lesions, and can be fatal

43 (Austin and Zhang, 2006; Korun, 2008; Mohamad et al., 2019). It can be caused by various species of Vibrio, such as Vibrio harveyi, Vibrio alginolyticus, Vibrio anguillarum, Vibrio ordalii, Vibrio 44 45 parahaemolyticus, and Vibrio vulnificus (Korun, 2008; Muniesa et al., 2020; Novriadi, 2016; Pujalte et al., 46 2003). Furthermore, the range of Vibrios spp. is expanding, posing an ongoing challenge in open-circuit 47 aquaculture systems supplied with water directly from the open sea (Baker-Austin et al., 2017). 48 Prophylactic measures, such as commercial vaccines, have been extensively developed and successfully 49 implemented to combat Vibrio anguillarum (Dadar et al., 2017). Now that the issue of V. anguillarum is 50 well managed, V. harveyi has recently emerged as a significant concern in seabass aquaculture (Abu Nor 51 et al., 2020; Islam et al., 2022; Vendramin et al., 2001).

52 Vibrio harveyi thrives in natural marine ecosystems. Its ability to act as a pathogen in a wide range of 53 cultivated hosts, such as crustaceans, molluscs, and fish is a concern notably for open-circuit aquaculture 54 farms (Austin and Zhang, 2006). Importantly, V. harveyi exhibits various states of existence, including free-55 swimming and within biofilms attached to both living and non-living surfaces, or engaged in symbiotic and 56 host-pathogen interactions (Montánchez and Kaberdin, 2020). The formation of biofilms on tank walls 57 can significantly heighten the risk of vibriosis outbreaks. Besides, outbreaks of vibriosis in seabass farms 58 exhibit pronounced seasonality, occurring more frequently when water temperatures reach their 59 maximum (Pujalte et al., 2003). Increasing temperatures, particularly with global warming, may boost the 60 abundance and possibly the virulence of viable V. harveyi in the surrounding environment, consequently 61 increasing the presence of V. harveyi and its damages in tanks. Actually, the occurrence of vibriosis 62 episodes remains unpredictable despite high temperatures, as highlighted by Mougin et al. (2021). In their 63 study, conducted in a small tank (75 m<sup>3</sup>), no instances of vibriosis occurred during the 7-month sampling 64 campaign, despite the clear impact of temperature on V. harveyi concentrations. The study was 65 conducted in a 1.3-metre-deep tank, which is not deep enough to observe the heterogeneity in 66 environmental parameters and their potential impact on V. harveyi and vibriosis development. This 67 underscores the need for further investigations into the relationships between environmental 68 parameters, V. harveyi concentrations dynamics, and vibriosis outbreaks. Additionally, to gain a more 69 comprehensive understanding of these relationships, studies should expand to larger tanks, where 70 environmental heterogeneity is more likely, allowing for a more nuanced examination of repartition 71 within the water column. Importantly, biofilms add an additional layer of complexity to the understanding 72 of vibriosis development. In biofilms, interspecific interactions are facilitated, and previous research 73 conducted by Roquigny et al. (2021) has already highlighted distinctions in microbial communities and 74 their temporal shifts in an aquaculture tank which may interact with V. harveyi. In this context, a first 75 question arises whether the temporal and spatial dynamics of these two forms of V. harveyi — free-76 swimming and within biofilms — are interconnected, and if one compartment of the tank acts as a 77 reservoir for each form. This uncertainty complicates the understanding of vibriosis onset, introducing 78 additional challenges in managing and mitigating the impact of V. harveyi in open sea environments. In 79 addition to the complexities posed by V. harveyi's presence in various states, vibriosis development is not 80 solely governed by host-pathogen interactions. Rather, it constitutes a multifaceted tripartite 81 relationship, influenced by environmental biotic and abiotic factors (Maulvault et al., 2018). These factors, 82 such as fish density, water temperature, salinity and pH, may collectively play a significant role in shaping 83 the dynamics of vibriosis outbreaks.

84 Thus, it is essential to comprehend V. harveyi's spatial and temporal distribution within aquaculture tanks, 85 as well as the environmental conditions that favour its proliferation and persistence. Therefore, through 86 a comprehensive investigation conducted from May to November 2022, encompassing sampling periods 87 before, during, and after episodes of vibriosis outbreaks, we sought to elucidate the spatial persistence 88 of V. harveyi in water and biofilms particularly during "heatwaves" or vibriosis outbreaks. This sampling 89 was done through both biofilm and water sampling, and environmental factors were measured alongside. 90 Besides, the conducted sampling was done at three different depths within the water column in a 4-metre 91 deep tank, to capture potential variations across vertical gradient. This study aimed to identify potential 92 environmental niches within these tanks where V. harveyi may thrive and persist, potentially leading to 93 vibriosis outbreaks. It could lay the groundwork necessary to effectively address its virulence and mitigate 94 the risk of vibriosis in aquaculture settings. Indeed, with escalating concerns over antibiotic resistance and 95 changing consumer preferences favouring sustainable practices, relying on antibiotic use is not a viable, 96 long-term solution. In this "One Health" context, aquaculture practitioners could solve this problem by 97 devising targeted antivirulence strategies that not only address the immediate challenges posed by *V*. 98 *harveyi*, but also align with the principles of sustainable and responsible aquaculture. The goal of this 99 research was to contribute to new strategies for effective management and regulation of vibriosis in 100 aquaculture facilities, based on informed preventive measures.

# 101 2. Material & methods

#### 102 2.1 Sampling site and rearing conditions

From May to November 2022, a sampling campaign was conducted at Aquanord-Ichtus, an aquaculture establishment situated in Gravelines (France), along the coast of the North Sea. This facility is strategically located near a nuclear power plant, which serves as a source of warm water for the flow-through tank systems via its reactor water cooling system. The facility benefits from thermoregulated water throughout the year, except in summer when the facility faces challenges in cooling the water tank more than environmental water.

109 The cubical aquaculture tank chosen is an outdoor facility with a volumetric capacity of approximately

110 975 cubic m, a height of 4 m, and a capacity of up to 68,000 sea bass. The average fish stock density within

the tank was 32 kg per cubic meter at the beginning of the campaign, and the average individual fish

112 weight was 517 g.

Fish are fed daily and oxygenation is carried out to maintain ideal dissolved oxygen (DO) levels. To overcome the absence of available vaccines against *V. harveyi*, Aquanord-Ichtus has started to create selfvaccines. This approach enables farmers to use oxytetracycline, an antibiotic, only in extreme cases when vibriosis -induced mortalities are exceptionally high and not effectively managed by other zootechnical measures. During vibriosis outbreaks and elevated fish mortality on the farm, the farmers administered

118 oxytetracycline treatment for 7–10 days, ensuring antibiotic coverage persists for about 4 weeks.

### 119 2.2 Sample collection and environmental parameters

120 To study the spatial and temporal dynamics of V. harveyi in biofilms, cylindrical concrete specimens (5 x 121 2.5 cm) constructed from the same material as the tank walls, were attached to a rope at three equidistant 122 depths from the surface of the tank (top: 0.3 m deep, middle: 2.15 m deep, and bottom: 4 m deep). At each sampling time, five replicates of concrete were sampled at each depth. To study planktonic V. 123 124 harveyi, alongside the concrete samples, 1 L of water in five replicates was also collected at each depth 125 using a submersible pump. This process yielded 16 independent sampling times, and samples taken from 126 16 ropes at the wall of the tank. Sampling apparatus was installed when the tank was in an empty state, 127 and the initial sampling was done one day after impoundment of the tank when the fish were added.

128 Biofilm and planktonic samples were collected every three weeks. Throughout the sampling campaign, 129 environmental seawater temperatures were carefully monitored, based on data provided by the MAREL 130 Carnot buoy consulted from May to November 2022 (MAREL Carnot, 2022). This buoy supplied daily 131 insights into the environmental temperatures of water used to partially supply the tanks. When seawater 132 temperatures exceeded 19°C for two consecutive nights, sampling events were scheduled within a span 133 of 4–5 days to anticipate the onset of vibriosis episodes. Given the sustained temperatures above 20°C 134 from mid-July to the end of August, a weekly sampling regimen was adopted, amounting to 7 samples 135 (Supplementary Table 1). At each sampling time, abiotic parameters such as water temperature, pH, 136 salinity, turbidity, DO, and conductivity were assessed within the tank at each layer, using an HI-9892 137 multi-parameter probe (Hanna Instruments, Leighton Buzzard, United Kingdom). Throughout the sampling campaign, the aquaculture facility supplied weekly data concerning biomass and mortality rates 138 139 for the studied rearing tank A total of 240 water samples and 240 biofilm samples were collected during 140 the sampling campaign.

### 141 2.3 Bacterial isolation

142 The experimental protocol, established for a previous study in Aquanord-Ichtus (Mougin et al., 2020), was 143 replicated with minor adaptations. Briefly, we gathered five replicates of 1 L seawater samples from 144 aquaculture tanks at different depths and sampling times. These samples were promptly chilled on ice 145 during transportation to the laboratory and processed within a 3-hour window following the sampling. 146 Filtration involved the sequential use of 0.45 µm and 0.22 µm-pore-size nitrocellulose filters (Grosseron 147 SAS, Couëron, France), with the deployment of multiple filters when necessary to prevent clogging due to 148 sample aggregation. Following this, the filters were immersed in a solution consisting of 30 mL of Luria-149 Bertani broth supplemented with 20 % NaCl (LBS) containing 20 % glycerol and stored at -80°C until further 150 analysis.

151 Concrete specimens were submerged at various depths: 0.3, 2.15, and 4 m. Post-collection, the specimens 152 were individually collected and preserved in sterile physiological water and put on ice during 153 transportation to the laboratory. They were immediately processed within a 2-hour timeframe upon 154 arrival. The biofilm collection protocol from concrete, following the Mougin et al. (2020) guidelines with 155 slight adaptations, used a systematic swabbing methodology after rinsing the blocks with sterile physiological water, targeting a specific surface area of 2.34 cm<sup>2</sup>. This process involved use of a sterile, 156 157 stainless steel jig and two consecutive sets of swabs (AB300250, Labomoderne, Gennevilliers, France). Subsequently, the swabs were immersed in LBS enriched with 20 % glycerol and stored at -80°C until 158 159 further analysis.

160 To begin the analytical phases, all samples were thawed and subjected to a 3-min vortexing process. After 161 centrifugation at 8,000 x g for 10 min, the resulting cell pellet was resuspended in physiological water. For 162 subsequent steps, the samples were divided into two subsamples, with one of them (1,800  $\mu$ L) to be used 163 for DNA extraction and *V. harveyi* quantification through real-time PCR, and the other (200  $\mu$ L) kept for 164 another study.

### 165 2.4 Quantification of *V. harveyi* using real-time PCR

The 1,800 µL subsamples underwent a centrifugation step at 8,000 x g for 10 min. Subsequently, a DNeasy 166 167 PowerBiofilm kit (QIAGEN, Hilden, Germany) was used for total DNA extraction from the cell pellet, 168 adhering to the manufacturer's instructions. This method was chosen based on the procedures described 169 by Mougin et al. (2020). This involved the use of V. harveyi-specific primers designed to target the mreb 170 gene, which is present in V. harveyi's genome as a single copy. The primers were synthesized by TIB 171 MOLBIOL Syntheselabor GmbH (Berlin, Germany). To establish standard curves for quantification, a series 172 of 10-fold dilutions was prepared using pure DNA from the collection strain V. harveyi LMG 4044. For each 173 individual assay, calibration curves were generated by plotting the mean threshold cycle (CT) values 174 derived from two replicate series against the logarithm of the respective dilution factors. To mitigate 175 potential interference from real-time PCR inhibitors, each extracted DNA template was analysed after 176 being diluted tenfold. Genome copy equivalent (GE) concentrations were calculated, taking the dilution 177 factor into consideration. All quantitative PCR reactions were conducted using a LightCycler® 480 178 instrument (Roche Diagnostics, Meylan, France). These assays included a positive control (V. harveyi LMG 179 4044 DNA), a negative control (V. campbellii CIP 70.67 DNA), and a no-template control (NTC) devoid of 180 DNA. To confirm the specificity of the fluorescent signals, melting curve analysis was performed. To ensure 181 the reliability of measurements, each extracted DNA template underwent two rounds of analysis.

### 182 2.5 Statistical analysis

183 Data normality was assessed using either the Shapiro–Wilk test or the Kruskal–Wallis test, depending on 184 the dataset distribution. Subsequently, the homogeneity of variances was evaluated using Bartlett's test. 185 To determine significant differences among variables across different layers, a one-way analysis of 186 variance (ANOVA) was conducted. Post-hoc comparisons were carried out using the least significant 187 difference (LSD) method with Bonferroni-adjusted p-values. In instances where the data did not meet the 188 assumption of normality, the Kruskal-Wallis test was employed as a non-parametric alternative for 189 comparing medians among different groups. Additionally, a principal component analysis (PCA) was 190 performed on environmental parameters (temperature, pH, salinity, DO, and turbidity), V. harveyi water 191 concentration, *V. harveyi* concrete concentration, and fish mortality. Before conducting PCA, the data 192 were standardised to ensure equal weighting of each variable. Relationships among variables were 193 explored using a correlation matrix, employing the Spearman method. Prior to this analysis, pairwise 194 linear relationships were assessed to ensure the absence of linearity. All statistical analyses, including 195 post-hoc procedures, were conducted using R version 4.3.2 and R Studio version 2021.09.0 (R Core Team, 196 2023; RStudio Team, 2020) with the "agricolae" (De Mendiburu and Yaseen, 2020), "factomineR" (Lê et 197 al., 2008), "factoextra" (Kassambara and Mundt, 2016), and "corrplot" (Wei and Simko, 2021) packages.

198 All graphics were drawn using the "ggplot2" packages (Wickham, 2016).

# 199 3. Results

#### 200 3.1 Water quality and fish mortality

201 The investigation involved a comprehensive examination of water parameters in the top, middle, and 202 bottom layers over the course of May to November 2022. Additionally, fish mortality data were collected 203 and analysed. These data are summarised in Table 1. No significant differences were found across the 204 water column for each sampling time for the recorded parameters, indicating homogeneity of 205 environmental conditions in the tank. Nevertheless, some parameters showed variations throughout the 206 sampling period. For instance, the mean temperature values ranged from a minimum of 16.70°C in 207 November to a maximum of 22.06 °C in August, demonstrating seasonal fluctuation. pH values gradually 208 decreased from an average of 8.07 in May to 6.87 in November. Values also showed increasing acidity 209 with depth in the water column. DO concentrations remained constant. Fish mortality percentages were 210 higher in July and highest in August when peak temperature values were recorded. Fish mortality episodes 211 in May were likely associated with stress induced by transfer of fish into the tank.

#### 212 3.2 Quantification of V. harveyi

Abundance of *V. harveyi* was determined using *mreb*-targeted real-time PCR across the different layers for all samples, as shown in (Fig. 1). The qPCR method has a specified threshold of detection in GE for each sample type: 2.24 GE.mL<sup>-1</sup> (0.35 log(GE.mL<sup>-1</sup>)) for water samples and 501.19 GE.cm<sup>-2</sup> (2.70 log(GE.cm<sup>-2</sup>)) for concrete samples. Despite the potential presence of *V. harveyi*, any concentration below this threshold was considered to be zero in the analysis (Supplementary Table 2). For a given sampling time, no significant differences were found trough the water column for each sample type.

219 In water samples, the presence of V. harveyi started to be detected from the beginning of the sampling 220 campaign in May, persisting consistently thereafter. A significant higher V. harveyi concentrations was 221 observed in August compared to other months (p < 0.05). Of note, V. harveyi abundance in the bottom 222 layer ranged from 0.52 to 3.36 log(GE.mL<sup>1</sup>) (Fig. 1), with a distinct peak in August when the temperature 223 of the water tank was at its highest. Moreover, abundances in the middle and top layers exhibited similar 224 profiles through the months, ranging from 0.60 to 2.43 log(GE.mL<sup>-1</sup>) and from 0.48 to 2.89 log(GE.mL<sup>-1</sup>) 225 respectively. Conversely, when oxytetracycline was used during episodes of mortality, a decrease in V. 226 harveyi concentrations was observed in water samples, allowing for concentration maintenance below 227 1.90 log(GE.mL<sup>-1</sup>), followed by a gradual increase towards the end of the antibiotic's predicted effect.

228 Many samples were found to be below the qPCR detection threshold for the concrete samples. The 229 estimated concentration of V. harveyi in biofilm samples did not exhibit a significant difference over time 230 and through the water column. Vibrio harveyi started to be detected in June and ceased by the end of 231 October for top and middle tank samples. Bottom samples were detected until the end of the sampling 232 campaign. Detected abundances in the bottom, middle and top layers of the tank ranged from 2.88 to 233 4.51 log(GE.cm<sup>-2</sup>), 3.02 to 4.60 log(GE.cm<sup>-2</sup>), and 2.98 to 4.93 log(GE.cm<sup>-2</sup>), respectively (Fig. 1). A rapid 234 increase in V. harveyi abundance was observed during high-temperature periods, particularly from August 235 to September, resulting in two peaks during these months associated with fish mortality in the tank (Table 236 1). Statistical analysis revealed significant variations in V. harveyi concentrations for all layers, with August 237 and September forming a cluster characterised by concentrations significantly higher than those observed for the rest of the months (p < 0.05). Moreover, during these episodes of mortality, oxytetracycline was 238 239 used and a resulting decrease in V. harveyi abundance was observed, enabling concentration maintenance to remain below 3.97 log(GE.cm<sup>-2</sup>). The abundance of *V. harveyi* gradually rose as the antibiotic's predicted

### 241 effect started to fade.

#### 242 3.3 Influence of environmental factors

243 PCA (Fig. 2) was combined with a correlation matrix (Fig. 3) to reveal the interactions of the parameters 244 measured in the aquaculture tank. Both dimensions of the PCA, Dim.1 and Dim.2, captured 64.40% of the 245 data variance, explaining 42.08% and 22.32% of total variance, respectively. Temperature, V. harveyi abundance in water ([Vh water]), and V. harveyi in biofilm form ([Vh concrete]), emerged as the primary 246 247 drivers shaping the Dim.1 variance landscape, while Dim.2 was predominantly influenced by DO and 248 turbidity. Strong positive correlations (0.88, p < 0.001) were found between V. harveyi abundance in water 249 and its abundance in biofilm within the tank, as highlighted in the correlation matrix, indicating that the 250 biofilm results align closely with those observed in the water. A strong positive correlation was observed 251 in the PCA between temperature and both V. harveyi abundance in water (0.891, p < 0.001) and in biofilm 252 (0.834, p < 0.001), but also, to a lesser extent, salinity (0.772, p < 0.001). Further confirming and extending 253 these findings, the correlation matrix highlighted significant positive correlations such as temperature with salinity (0.84, p < 0.001), V. harveyi concentration in water (0.63, p < 0.001), concrete (0.55, p < 0.001) 254 255 0.001), and fish mortality (0.50, p < 0.001). In contrast, pH exhibited a negative correlation with V. harveyi 256 abundance in water (-0.43, p < 0.001), indicating an inverse relationship. On the other hand, Dim.2 257 revealed significant correlations with DO (0.798, p < 0.001), pH (0.566, p < 0.001), fish mortality (0.496, p258 < 0.001), and turbidity (-0.741, p < 0.001). Additionally, the correlation matrix revealed strong negative 259 correlations between DO and turbidity (-0.40, p = 0.0054), and positive correlation with fish mortality 260 (0.34, p = 0.0196).

## 261 4. Discussion

#### 262 4.1 Temporal and spatial study of *V. harveyi* in aquaculture

Our investigation examined the temporal and spatial distribution of V. harveyi throughout a sampling 263 264 campaign spanning May to November 2022 on a farm in northern France. This farm has experienced 265 recurring vibriosis episodes attributed to V. harveyi. A previous study conducted in a "small tank" (75 m<sup>3</sup>, 266 1.3 m deep), revealed the consistent presence and persistence of V. harveyi in both planktonic form and 267 biofilms throughout the year (Mougin et al., 2021). This previous study was restricted by the absence of 268 vibriosis outbreaks during the campaign, but also by the small tank size, which prevented in-depth 269 exploration of variations in environmental conditions and their impact on V. harveyi. To better understand 270 the dynamics of V. harveyi through the water column, we implemented sampling campaign in a 4-metre-271 deep tank at three different depths. We systematically monitored environmental parameters to find 272 potential explanations for the observed variations. Our study sets itself apart by adopting a 273 comprehensive sampling approach over an extended time frame that included the periods before, during, 274 and after two episodes of vibriosis. Additionally, during our sampling campaign, the two vibriosis 275 outbreaks forced the farmers to use antibiotics. Consequently, this study offers insights into antibiotic 276 effects on V. harveyi, considering both planktonic and biofilm forms. This campaign also sheds light on the 277 omnipresence of V. harveyi within the aquaculture tank. From the very onset, brought in with the initial 278 filling of the tank with water, V. harveyi demonstrated its adaptability, swiftly establishing its presence in 279 the tank environment. Vibrio harveyi was present and quantifiable during the whole sampling campaign 280 in planktonic form in the water, and during warmer months in biofilms. Its abundance in biofilms 281 potentially increased due to persistence once established on the tank wall, a finding in line with the 282 previous sampling campaign on the same aquaculture farm in 2018 (Mougin et al., 2021). A distinctive 283 characteristic of Vibrio, including V. harveyi, is their ability to adapt to changing conditions. This 284 adaptability is underscored by their capacity to transition from free-swimming cells to "swarmer cells" 285 within biofilms (Khan et al., 2020; McCarter, 1999). Furthermore, during the warmer months, it is known 286 that the abundance of other coexisting bacterial families within biofilms such as Rhodobacteraceae 287 decreases, providing an environment that enables the Vibrionaceae to thrive in the microbial community 288 structure (Bourne et al., 2004; Roquigny et al., 2021)

#### 289 4.2 Abundance of *V. harveyi* in the tank

290 In comparison to Mougin et al. (2021) our study observed higher concentrations of V. harveyi in our water 291 tank. They found, actually, a maximum concentration of 9.51×10<sup>1</sup> GE.mL<sup>-1</sup> in rearing tanks, while a 292 maximum concentration of 2.54 x 10<sup>3</sup> GE.ml<sup>-1</sup> was found in our water tank. This contrast could be due to 293 differences in environmental conditions, but also due to different tank sizes and fish. In our study, which 294 aimed to investigate the water column, all measured abiotic parameters in the water column exhibited 295 clear stability, a characteristic likely shaped by the conditions of fish farming. These conditions, which 296 ensure efficient water circulation, additionally play a role in homogenising tank parameters overall. The 297 concentrations of V. harveyi reflected the same stability through the water column. Similar observations 298 regarding Vibrio spp. concentrations in the water column were found in a natural oyster farming site (Cruz 299 et al., 2020). The only significant difference observed by Cruz et al. (2020) appeared for some oysters 300 collected close to the seafloor where sediment accumulation was high. Even though our tank water 301 appeared to be homogeneous, organic matter accumulated at the floor of the tank. This phenomenon is 302 caused by fewer fish activities at the bottom, resulting in less associated movement. It can also be 303 attributed to the sedimentation and accumulation of nutrient-rich sediments consisting of food scraps 304 and fish faeces. Moreover, sediments are known to serve as a source of nutrients for V. harveyi, which 305 not only sediment themselves in the tank, but are also able to absorb dissolved organic matter (Neogi et 306 al., 2018). Sediments also provide an additional surface for colonisation, protecting the bacteria from 307 biotic and abiotic stresses, and enabling their persistence in the tank (Pianetti et al., 2004; Vezzulli et al., 308 2002). This hypothesis regarding sediment dynamics and V. harveyi colonization could be tested by 309 conducting further sampling after emptying the tank, which would provide insights into the role of 310 sediments in V. harveyi dynamics in aquaculture settings.

#### 4.3 Environmental factors influencing *V. harveyi* distribution

312 Even though V. harveyi concentrations were spatially homogeneous throughout the water column, 313 temporal variations were observed through the 7-month sampling campaign. Relationships between 314 bacterial concentrations and measured environmental parameters were investigated through PCA (Fig. 2) 315 and a correlation matrix (Fig. 3). Interestingly, temperature emerged as the foremost factor influencing 316 the abundance of V. harveyi, as abundance was consistently noted to be more favourable as temperatures 317 rose above 20 °C. Given that V. harveyi was introduced into the tank water from the sea, its persistence 318 within the tank is facilitated by favourable tank conditions, enabling it to form biofilms on the tank walls 319 and promoting its growth. Additionally, extensive studies have demonstrated the crucial influence of 320 temperature on the presence of V. harveyi in natural aquatic systems, displaying a clear seasonal cycle, 321 particularly during the summer when temperatures exceed 25 °C (Chimetto et al., 2008; Zhou et al., 2012). 322 This seasonality was also observed in our study, as V. harveyi exhibited higher abundance during the 323 warmest months, when temperatures reached up to 22 °C. However, it is important to note that vibriosis 324 does not necessarily occur when temperature reach 22 °C, suggesting that additional factors may 325 influence its occurrence. Additionally, the correlation matrix (Fig. 3) suggests that salinity and pH, which 326 are predicted to change with overall ocean temperatures (Gulev et al., 2021), also positively affected the 327 presence of V. harveyi in the aquaculture tank. These findings are consistent with prior researches on the 328 effects of environmental parameters on Vibrio species in natural and in aquaculture settings, where a 329 positive correlation between salinity and temperature with the frequency of detection of Vibrio species 330 has been reported (Abioye et al., 2021; Scro et al., 2022). DO concentration is reported to have a negative 331 correlation with temperature, given that increased water temperatures lead to decreased DO levels 332 (Pfeffer et al., 2003). However, in our case, DO levels were managed continuously by the farmers, which 333 explains the absence of any observable effect on the abundance of V. harveyi. A second negative 334 correlation was observed with pH values. Although V. harveyi is known to tolerate wide ranges of pH levels 335 (i.e., 5.5 to 9.0), the correlation matrix in our study shows that pH is negatively correlated with both V. 336 harveyi abundance in water and salinity (Prayitno and Latchford, 1995). This parameter cannot be fully 337 controlled in aquaculture, adding another challenge to vibriosis management in tanks. In addition, global 338 warming and ocean acidification may alter the microbial balance, favouring the presence of Vibrio, while 339 diminishing populations of other bacterial genera (Montánchez and Kaberdin, 2020). All these observations highlight the complexity to identify the multiparameter ideal *Vibrio* environment; varying

based on geographic location and environmental conditions (Trinanes and Martinez-Urtaza, 2021).

## **342** 4.4 Challenges and considerations in managing vibriosis outbreaks

343 The repeated occurrences and unpredictability of vibriosis outbreaks associated with V. harveyi present a 344 significant challenge to the seabass aquaculture sector, as highlighted by Vendramin et al. (2001) and 345 Firmino et al. (2019). The issue becomes particularly problematic in open-system aquaculture, since the 346 prevalence of this pathogen in fish closely aligns with its abundance in the surrounding environment (Kim 347 and Lee, 2017), but also because water quality cannot be fully controlled. Moreover, in the context of 348 climate change, there has been a temperature increase of about 0.88 °C in sea and ocean surface temperatures over the last century (Gulev et al., 2021). This temperature rise is known to affect the 349 350 distribution and abundance of marine organisms (IPCC, 2022), including Vibrio, which has seen its 351 abundance increase sharply in the North Sea over the period 1958–2011 (Vezzulli et al., 2002). According 352 to the Intergovernmental Panel on Climate Change, the ocean is projected to continue warming, leading to long-term impacts on ocean ecosystems and human societies (IPCC, 2022). Open aquaculture systems 353 354 may therefore need to address an increase in vibriosis episodes. This heightened risk could require increased antibiotic use, posing risks to fish health, environmental integrity, and consumer safety. In 355 356 response to these challenges, it is imperative to develop specific and effective solutions. Antibiotics, as 357 suggested by our findings, fall short as a sustainable solution, since despite their use by farmers, the 358 concentrations of V. harveyi in water and biofilms quickly rebound. Within a three-month period, farmers 359 had to resort to antibiotics twice to avoid massive fish losses (Fig. 2). Given the escalating importance of addressing the Vibrio challenge in response to global warming, it is crucial to adopt a comprehensive 360 361 approach to understand the virulence mechanisms used by V. harveyi. As an opportunistic pathogen, V. 362 harveyi thrives in warmer temperatures, requiring proactive measures in response to changing climate 363 conditions that could now favour its growth in previously cooler waters. Addressing the core issue involves 364 reducing bacterial virulence through antivirulence therapies (Clatworthy et al., 2007; D'Angelo et al., 365 2018). Environmental effects impact not only to behaviour of the bacterium, but also to its area of 366 distribution, which is bound to evolve. For some geographic areas, including the one in the present study 367 and more generally temperate waters, such changes may pose challenges regarding Vibrio expansion 368 (Trinanes and Martinez-Urtaza, 2021). Moreover, they may even compromise host immunity, thereby 369 enhancing the bacterium's potential to increase mortality. Understanding these multifaceted interactions 370 becomes a keystone in formulating nuanced and effective strategies aimed at mitigating the impact of 371 potential vibriosis outbreaks.

# 372 5. Conclusion

373 In conclusion, this study highlighted the challenges of managing V. harveyi outbreaks in aquaculture 374 systems. The persistence of V. harveyi pose major challenges, particularly during warmer months. The 375 high adaptability of Vibrio and particularly V. harveyi, implies its considerable permissiveness, rendering 376 the task of attempting to control the entire environment to prevent vibriosis highly challenging, especially 377 in the context of global warming and ocean acidification (Montánchez and Kaberdin, 2020). Additionally, 378 significant correlations were found between V. harveyi concentrations and various environmental 379 parameters, highlighting the complex dynamics influencing its distribution. While temperature emerged 380 as a primary influencer, the absence of vibriosis outbreaks every summer suggests the influence of 381 additional parameters such as salinity and pH. Furthermore, in the context of escalating Vibrio abundance 382 linked to rising global ocean temperatures, concerns arise regarding the heightened pathogenic potential 383 of certain Vibrio strains, potentially leading to an increased incidence of diseases in the marine 384 environment. This poses a real challenge for open aquaculture systems which are specifically impacted by 385 these temperature increases as they are unable to regulate water temperature during the hottest months. 386 In navigating these intricate ecological challenges, the integration of a "One Health" framework becomes 387 imperative, emphasising the interconnectedness of animal, environmental, and human health (McEwen 388 and Collignon, 2018; Zinsstag et al., 2011). Understanding the virulence mechanisms of V. harveyi is crucial 389 for developing effective countermeasures and adaptive strategies that align with changing ecological 390 systems. This emphasizes the need to explore its virulence genes and their responses to environmental factors like temperature, which are expected to evolve in the future. These insights are crucial for guiding

392 future research and developing strategies for disease management and sustainable aquaculture practices.

# 393 6. Acknowledgements

The authors would like to thank Aquanord-Ichtus and especially Guillaume Tielie, Anabelle Duhamel, and Céline Doyen. The authors also thank Lorédane Clause for technical support. Alix Da Fonseca Ferreira would like to thank the PMCO council and Hauts-de-France regional council for their financial support of her PhD studies. This work was funded by the French government, Ifremer, and the Hauts-de-France region in the framework of the CPER 2021–2027 IDEAL project, and by IFSEA Graduate School (which originates from the French National Research Agency under the Investments for the Future programme ANR-21-EXES-00 11).

# 401 7. References

- 402Abioye, O.E., Osunla, A.C., Okoh, A.I., 2021. Molecular detection and distribution of six medically403important Vibrio spp. in selected freshwater and brackish water resources in eastern cape404province, south africa. Frontiers in Microbiology 12. https://doi.org/10.3389/fmicb.2021.617703
- Abu Nor, N., Zamri-Saad, M., Md Yasin, I.-S., Salleh, A., Mustaffa-Kamal, F., Matori, M.F., Azmai, M.N.A.,
  2020. Efficacy of Whole Cell Inactivated *Vibrio harveyi* Vaccine against vibriosis in a Marine Red
  Hybrid Tilapia (*Oreochromis niloticus × O. mossambicus*) Model. Vaccines (Basel) 8, 734.
  https://doi.org/10.3390/vaccines8040734
- Ahmed, N., Thompson, S., Glaser, M., 2019. Global Aquaculture Productivity, Environmental
   Sustainability, and Climate Change Adaptability. Environmental Management 63, 159–172.
   https://doi.org/10.1007/s00267-018-1117-3
- ANSES, 2019. Fiche de description de danger biologique transmissible par les aliments : Vibrions entéropathogènes : *Vibrio parahaemolyticus, Vibrio cholerae* non-O1/ non-O139 et V*ibrio vulnificus* - Décembre 2019 [WWW Document]. Anses - Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail. URL https://www.anses.fr/fr/content/fiche-dedescription-de-danger-biologique-transmissible-par-les-aliments-vibrions (accessed 11.14.22).
- Austin, B., Zhang, X.-H., 2006. *Vibrio harveyi*: a significant pathogen of marine vertebrates and invertebrates. Letters in Applied Microbiology 43, 119–124. https://doi.org/10.1111/j.1472-765X.2006.01989.x
- Baker-Austin, C., Trinanes, J., Gonzalez-Escalona, N., Martinez-Urtaza, J., 2017. Non-Cholera Vibrios: The
   Microbial Barometer of Climate Change. Trends Microbiol 25, 76–84.
   https://doi.org/10.1016/j.tim.2016.09.008
- Bourne, D.G., Young, N., Webster, N., Payne, M., Salmon, M., Demel, S., Hall, M., 2004. Microbial
  community dynamics in a larval aquaculture system of the tropical rock lobster, *Panulirus ornatus*.
  Aquaculture 242, 31–51. https://doi.org/10.1016/j.aquaculture.2004.08.047
- Chimetto, L.A., Brocchi, M., Thompson, C.C., Martins, R.C.R., Ramos, H.R., Thompson, F.L., 2008. Vibrios
   dominate as culturable nitrogen-fixing bacteria of the Brazilian coral *Mussismilia hispida*.
   Systematic and Applied Microbiology 31, 312–319. https://doi.org/10.1016/j.syapm.2008.06.001
- Clatworthy, A.E., Pierson, E., Hung, D.T., 2007. Targeting virulence: a new paradigm for antimicrobial
   therapy. Nature chemical biology 3, 541–8. https://doi.org/10.1038/nchembio.2007.24
- 431 Cruz, C.D., Fletcher, G.C., Paturi, G., Hedderley, D.I., 2020. Influence of farming methods and seawater
   432 depth on *Vibrio* species in New Zealand Pacific oysters. International Journal of Food Microbiology
   433 325, 108644. https://doi.org/10.1016/j.ijfoodmicro.2020.108644
- Dadar, M., Dhama, K., Vakharia, V.N., Hoseinifar, S.H., Karthik, K., Tiwari, R., Khandia, R., Munjal, A.,
  Salgado-Miranda, C., Joshi, S.K., 2017. Advances in Aquaculture Vaccines Against Fish Pathogens:
  Global Status and Current Trends. Reviews in Fisheries Science & Aquaculture 25, 184–217.
  https://doi.org/10.1080/23308249.2016.1261277
- D'Angelo, F., Baldelli, V., Halliday, N., Pantalone, P., Polticelli, F., Fiscarelli, E., Williams, P., Visca, P., Leoni,
   L., Rampioni, G., 2018. Identification of FDA-Approved Drugs as Antivirulence Agents Targeting

- 440 the pqs Quorum-Sensing System of *Pseudomonas aeruginosa*. Antimicrobial Agents and 441 Chemotherapy 62, e01296-18. https://doi.org/10.1128/AAC.01296-18
- 442 De Mendiburu, F., Yaseen, M., 2020. agricolae: Statistical Procedures for Agricultural Research.
- 443 EUMOFA, 2022. The EU fish market: 2022 edition. Publications Office, LU.
- Firmino, J., Furones, M.D., Andree, K.B., Sarasquete, C., Ortiz-Delgado, J.B., Asencio-Alcudia, G., Gisbert,
  E., 2019. Contrasting outcomes of *Vibrio harveyi* pathogenicity in gilthead seabream, *Sparus aurata* and European seabass, *Dicentrachus labrax*. Aquaculture 511, 734210.
  https://doi.org/10.1016/j.aquaculture.2019.734210
- Gulev, S.K., Thorne, P.W., Ahn, J., Dentener, F.J., Domingues, C.M., Gerland, S., Gong, D., Kaufman, D.S.,
   Nnamchi, H.C., Quaas, J., 2021. Changing state of the climate system. Climate change 287–422.
- IPCC, 2022. Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group
   II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC Sixth
   Assessment Report.
- Islam, S.I., Mou, M.J., Sanjida, S., Tariq, M., Nasir, S., Mahfuj, S., 2022. Designing a novel mRNA vaccine
   against *Vibrio harveyi* infection in fish: an immunoinformatics approach. Genomics Inform 20,
   e11. https://doi.org/10.5808/gi.21065
- 456 Kassambara, A., Mundt, F., 2016. factoextra: Extract and Visualize the Results of Multivariate Data 457 Analyses.
- Khan, F., Tabassum, N., Anand, R., Kim, Y.-M., 2020. Motility of *Vibrio* spp.: regulation and controlling
  strategies. Appl Microbiol Biotechnol 104, 8187–8208. https://doi.org/10.1007/s00253-02010794-7
- Kim, J.Y., Lee, J.-L., 2017. Correlation of Total Bacterial and *Vibrio* spp. Populations between Fish and
   Water in the Aquaculture System. Frontiers in Marine Science 4.
- Korun, J., 2008. Marine *Vibrios* associated with diseased sea bass (*Dicentrarchus labrax*) in Turkey. Journal
   of Fisheriessciences.com. https://doi.org/10.3153/jfscom.2008007
- Lê, S., Josse, J., Husson, F., 2008. FactoMineR: A Package for Multivariate Analysis. Journal of Statistical
   Software 25, 1--18. https://doi.org/10.18637/jss.v025.i01
- MAREL Carnot, 2023. High Frequency measurement of the coastal environment in the eastern English
   Channel. Data from MAREL CARNOT COAST-HF (Coastal ocean observing system High
   frequency) monitoring programme within the Research Infrastructure ILICO.
   https://doi.org/10.17882/39754
- Maulvault, A.L., Camacho, C., Barbosa, V., Alves, R., Anacleto, P., Fogaça, F., Kwadijk, C., Kotterman, M.,
  Cunha, S.C., Fernandes, J.O., Rasmussen, R.R., Sloth, J.J., Aznar-Alemany, O., Eljarrat, E., Barceló,
  D., Marques, A., 2018. Assessing the effects of seawater temperature and pH on the
  bioaccumulation of emerging chemical contaminants in marine bivalves. Environmental Research
  161, 236–247. https://doi.org/10.1016/j.envres.2017.11.017
- 476 McCarter, L., 1999. The multiple identities of *Vibrio parahaemolyticus*. J Mol Microbiol Biotechnol 1, 51–
  477 57.
- 478 McEwen, S.A., Collignon, P.J., 2018. Antimicrobial Resistance: a One Health Perspective. Microbiology
   479 Spectrum 6, 6.2.10. https://doi.org/10.1128/microbiolspec.ARBA-0009-2017
- Mohamad, N., Amal, M.N.A., Saad, M.Z., Yasin, I.S.M., Zulkiply, N.A., Mustafa, M., Nasruddin, N.S., 2019.
   Virulence-associated genes and antibiotic resistance patterns of *Vibrio* spp. isolated from cultured
   marine fishes in Malaysia. BMC veterinary research 15, 176. https://doi.org/10.1186/s12917-019 1907-8
- Montánchez, I., Kaberdin, V.R., 2020. *Vibrio harveyi*: A brief survey of general characteristics and recent
   epidemiological traits associated with climate change. Marine Environmental Research 154,
   104850. https://doi.org/10.1016/j.marenvres.2019.104850
- Mougin, J., Roquigny, R., Flahaut, C., Bonnin-Jusserand, M., Grard, T., Le Bris, C., 2021. Abundance and
  spatial patterns over time of Vibrionaceae and *Vibrio harveyi* in water and biofilm from a seabass
  aquaculture facility. Aquaculture 542, 736862.
  https://doi.org/10.1016/j.aquaculture.2021.736862
- 491 Mougin, J., Roquigny, R., Travers, M.-A., Grard, T., Bonnin-Jusserand, M., Le Bris, C., 2020. Development 492 of a mreB-targeted real-time PCR method for the quantitative detection of *Vibrio harveyi* in

- 493seawater and biofilm from aquaculture systems. Aquaculture 525, 735337.494https://doi.org/10.1016/j.aquaculture.2020.735337
- Muniesa, A., Basurco, B., Aguilera, C., Furones, D., Reverté, C., Sanjuan-Vilaplana, A., Jansen, M.D., Brun,
  E., Tavornpanich, S., 2020. Mapping the knowledge of the main diseases affecting sea bass and
  sea bream in Mediterranean. Transboundary and Emerging Diseases 67, 1089–1100.
  https://doi.org/10.1111/tbed.13482
- 499 Neogi, S.B., Lara, R., Alam, M., Harder, J., Yamasaki, S., Colwell, R.R., 2018. Environmental and
   500 hydroclimatic factors influencing *Vibrio* populations in the estuarine zone of the Bengal delta.
   501 Environ Monit Assess 190, 565. https://doi.org/10.1007/s10661-018-6925-7
- 502
   Novriadi,
   R.,
   2016.
   Vibriosis
   in
   Aquaculture.
   Omniakuatika
   12,
   1–12.

   503
   https://doi.org/10.20884/1.oa.2016.12.1.24

   12,
   1-12.

   <t
- Pfeffer, C.S., Hite, M.F., Oliver, J.D., 2003. Ecology of *Vibrio vulnificus* in Estuarine Waters of Eastern North
   Carolina. Applied and Environmental Microbiology 69, 3526–3531.
   https://doi.org/10.1128/AEM.69.6.3526-3531.2003
- Pianetti, A., Bruscolini, F., Sabatini, L., Colantoni, P., 2004. Microbial characteristics of marine sediments
   in bathing area along Pesaro-Gabicce coast (Italy): a preliminary study. Journal of Applied
   Microbiology 97, 682–689. https://doi.org/10.1111/j.1365-2672.2004.02352.x
- 510 Prayitno, S.B., Latchford, J.W., 1995. Experimental infections of crustaceans with luminous bacteria related to Photobacterium and Vibrio. Effect of salinity and pH on infectiosity. Aquaculture, Fifth 511 512 International Colloguium on Pathology in Marine Aquaculture 132, 105–112. 513 https://doi.org/10.1016/0044-8486(94)00374-W
- Pujalte, M.J., Sitjà-Bobadilla, A., Macián, M.C., Belloch, C., Álvarez-Pellitero, P., Pérez-Sánchez, J., Uruburu,
   F., Garay, E., 2003. Virulence and Molecular Typing of *Vibrio harveyi* Strains Isolated from Cultured
   Dentex, Gilthead Sea Bream and European Sea Bass. Systematic and Applied Microbiology 26,
   284–292. https://doi.org/10.1078/072320203322346146
- 518 R Core Team, 2023. R: A Language and Environment for Statistical Computing.
- Roquigny, R., Mougin, J., Le Bris, C., Bonnin-Jusserand, M., Doyen, P., Grard, T., 2021. Characterization of
   the marine aquaculture microbiome: A seasonal survey in a seabass farm. Aquaculture 531,
   735987. https://doi.org/10.1016/j.aquaculture.2020.735987
- 522 RStudio Team, 2020. RStudio: Integrated Development for R. RStudio.
- Scro, A.K., Westphalen, J., Kite-Powell, H.L., Brawley, J.W., Smolowitz, R.M., 2022. The effect of off-bottom
   versus on-bottom oyster culture on total and pathogenic *Vibrio* spp. abundances in oyster tissue,
   water and sediment samples. International Journal of Food Microbiology 379, 109870.
   https://doi.org/10.1016/j.ijfoodmicro.2022.109870
- Trinanes, J., Martinez-Urtaza, J., 2021. Future scenarios of risk of *Vibrio* infections in a warming planet: a
   global mapping study. The Lancet Planetary Health 5, e426–e435. https://doi.org/10.1016/S2542 5196(21)00169-8
- Vendramin, N., Zrncic, S., Padrós, F., Oraic, D., Breton, A., Zarza, C., Olesen, N., 2001. Fish health in
   Mediterranean Aquaculture, past mistakes and future challenges. Bulletin- European Association
   of Fish Pathologists 36, 38–45.
- Vezzulli, L., Chelossi, E., Riccardi, G., Fabiano, M., 2002. Bacterial community structure and activity in fish
   farm sediments of the Ligurian sea (Western Mediterranean). Aquaculture International 10, 123–
   141. https://doi.org/10.1023/A:1021365829687
- 536 Wei, T., Simko, V., 2021. R package "corrplot": Visualization of a Correlation Matrix.
- 537 Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
- Zhou, J., Fang, W., Yang, X., Zhou, S., Hu, L., Li, X., Qi, X., Su, H., Xie, L., 2012. A nonluminescent and highly
   virulent *Vibrio harveyi* strain is associated with "bacterial white tail disease" of *Litopenaeus* vannamei shrimp. PLoS One 7, e29961. https://doi.org/10.1371/journal.pone.0029961
- Zinsstag, J., Schelling, E., Waltner-Toews, D., Tanner, M., 2011. From "One medicine" to "One health" and
   systemic approaches to health and well-being. Preventive Veterinary Medicine 101, 148–156.
   https://doi.org/10.1016/j.prevetmed.2010.07.003
- 544

545 *Figure 1:* Biofilm concentrations of *V. harveyi* in log genome copy equivalents (GE).cm<sup>-2</sup> and *V. harveyi* concentrations in log

546 GE.mL<sup>-1</sup> according to tank depth and time. The coloured lines correspond to the averages of the 5 replicates, which are 547 represented by the shapes. Antibiotic treatments with oxytetracycline are indicated in dark green, and their period of effect is

star represented by the snapes. Antibiotic treatments with oxytetracycline are indicated in dark green, and their period of effectsstarted in light green. The dashed line marks the qPCR detection threshold for *V. harveyi* quantification, delineating the limit

549 above which presence is reliably detected.

Figure 2: Circle of correlation of the PCA for parameters including *V. harveyi* in water ([*Vh* water]), *V. harveyi* in biofilms ([*Vh* concrete]), dissolved oxygen, temperature, salinity, turbidity, pH, and fish mortality (Mortality). The contribution of each variable indicates its importance in explaining the overall variance in the dataset.

Figure 3: Correlation matrix of tank environmental parameters including *V. harveyi* concentrations in water ([*Vh* Water]), *V. harveyi* in biofilms ([*Vh* Concrete]), dissolved oxygen, temperature, salinity, turbidity, pH, and fish mortality (Mortality). The size
 of the circles corresponds to the strength of correlation, with larger circles indicating stronger correlations and small ones
 indicating weak correlations. Only statistically significant correlations (*p* < 0.05) between the parameters are shown.</li>

557 *Table 1:* Mean values ± standard deviations of parameters recorded from May to November 2022 in aquaculture facilities,
 558 categorised by the top, middle (mid), and bottom (bot) layers of the rearing tank. The 'n' signifies the number of sampling times
 559 averaged. For each sampling time, the average was calculated from 10 probe measurements.

		Sampling month						
		<b>May</b> (n = 2)	<b>June</b> (n = 2)	<b>July</b> (n = 3)	August (n = 4)	September (n = 2)	October (n = 2)	November (n = 1)
(c°)	Тор	18.29 ± 0.77	19.06 ± 0.30	20.89 ± 0.28	22.06 ± 0.58	20.69 ± 0.03	17.82 ± 0.36	16.71 ± 0
perature	Mid	18.29 ± 0.74	19.06 ± 0.30	20.89 ± 0.28	22.05 ± 0.57	20.71 ± 0.05	17.82 ± 0.36	16.70 ± 0
Tem	Bot	18.29 ± 0.74	19.06 ± 0.30	20.89 ± 0.28	22.06 ± 0.57	20.71 ± 0.04	17.82 ± 0.36	16.70±0
	Top	8.07 ± 0.71	7.53 ± 0.16	7.37 ± 0.22	7.29 ± 0.04	7.39 ± 0.06	7.17 ± 0.19	7.06 ± 0
Hq	Mid	8.06 ± 0.70	7.55 ± 0.06	7.31 ± 0.24	7.23 ± 0.07	7.33 ± 0.03	7.12 ± 0.17	7.03 ± 0
	Bot	7.95 ± 0.72	7.51 ± 0.02	7.25 ± 0.23	7.18 ± 0.07	7.25 ± 0.08	7.02 ± 0.16	6.87 ± 0
inity (psu)	Top	33.83 ± 0.68	34.75 ± 0.16	35.14 ± 0.16	35.07 ± 0.16	34.97 ± 0.11	34.09 ± 0.05	33.08 ± 0
	Mid	33.40 ± 1.34	34.74 ± 0.18	35.17 ± 0.18	35.16 ± 0.07	34.91 ± 0.05	34.00 ± 0.18	31.99 ± 0
Sal	Bot	33.45 ± 1.34	34.63 ± 0.27	35.12 ± 0.15	35.18 ± 0.10	34.83 ± 0.20	33.81 ± 0.30	32.67 ± 0
ygen	Top	5.27 ± 0.15	4.13 ± 0.04	3.20 ± 0.01	4.70 ± 0.21	5.46 ± 0.02	3.47 ± 0.06	6.47 ± 0.15
lved ox (mg.L <sup>-1</sup> )	Mid	5.45 ± 0.03	5.19 ± 0.02	3.39 ± 0.01	3.63 ± 0.02	5.83 ± 0.09	3.46 ± 0.04	3.64 ± 0.26
Disso	Bot	5.74 ± 0.03	5.49 ± 0.03	3.44 ± 0.02	4.04 ± 0.03	7.30 ± 0.43	3.82 ± 0.05	3.65 ± 0.11
NU)	Тор	4.51 ± 1.02	5.02 ± 0.07	4.90 ± 1.44	2.84 ± 0.82	6.89 ± 0.52	7.15 ± 4.95	3.30 ± 0
oidity (F	Mid	4.26 ± 1.32	3.40 ± 1.45	4.17 ± 0.75	3.86 ± 0.83	4.64 ± 1.65	5.56 ± 2.36	6.67 ± 0
Turb	Bot	3.99 ± 0.47	3.99 ± 0.77	6.41 ± 2.73	3.83 ± 0.80	6.39 ± 3.13	5.35 ± 1.77	5.29 ± 0

-								
	Fish Mortality (%)	1.26	0.98	1.31	4.57	0.90	0.06	0.01
560	_							. O